

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 1  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %	
0101	58	*	*	*	*	*	*	
0102	159	95.0	2.5	2.5	0.0	0.0	0.0	
0103	168	99.4	0.6	0.0	0.0	0.0	0.0	
0104	58	*	*	*	*	*	*	
0250	64	*	*	*	*	*	*	
0251	41	*	*	*	*	*	*	
0252	55	*	*	*	*	*	*	
0253	52	*	*	*	*	*	*	
0401	195	97.4	0.0	0.0	0.0	0.0	2.6	
0402	172	90.7	0.6	1.2	0.0	0.0	7.6	
0403	60	*	*	*	*	*	*	
0404	55	*	*	*	*	*	*	
0501	203	97.5	1.0	1.5	0.0	0.0	0.0	
0502	2	*	*	*	*	*	*	
0503	68	*	*	*	*	*	*	
0504	239	89.1	0.4	0.8	3.8	0.4	5.4	
0505	155	85.2	1.3	3.2	0.6	0.0	9.7	
0506	74	*	*	*	*	*	*	
0650	382	89.5	2.4	3.1	1.3	1.3	2.4	
0651	252	97.6	0.8	1.2	0.0	0.4	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 2  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
0652	136	97.1	0.0	1.5	0.0	0.0	1.5
0653	147	98.0	0.0	0.7	0.0	0.0	1.4
0654	128	98.4	0.8	0.0	0.0	0.0	0.8
0655	130	100.0	0.0	0.0	0.0	0.0	0.0
0657	42	*	*	*	*	*	*
0701	173	91.3	0.6	1.2	0.6	5.8	0.6
0702	38	*	*	*	*	*	*
0703	94	*	*	*	*	*	*
0704	32	*	*	*	*	*	*
0705	59	*	*	*	*	*	*
0850	399	89.2	5.5	3.0	0.0	0.8	1.5
0851	337	95.5	0.0	3.9	0.0	0.0	0.6
0901	675	86.2	3.3	8.0	0.3	0.6	1.6
0902	21	*	*	*	*	*	*
1051	540	90.0	1.1	5.9	1.5	0.4	1.1
1101	724	37.7	55.4	4.0	0.4	1.7	0.8
1102	336	80.7	10.4	6.0	0.6	0.9	1.5
1106	49	*	*	*	*	*	*
1250	387	88.9	0.3	9.0	0.0	0.8	1.0
1251	283	97.9	1.1	1.1	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 3  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
1252	293	64.8	1.4	33.1	0.0	0.0	0.7
1253	278	86.3	10.4	2.9	0.0	0.0	0.4
1350	905	67.1	22.1	9.2	0.2	0.6	0.9
1351	56	*	*	*	*	*	*
1352	153	63.4	31.4	4.6	0.0	0.0	0.7
1401	110	94.5	0.9	0.9	0.0	0.0	3.6
1402	61	*	*	*	*	*	*
1403	8	*	*	*	*	*	*
1404	8	*	*	*	*	*	*
1501	423	55.6	7.8	35.7	0.2	0.0	0.7
1502	45	*	*	*	*	*	*
1503	404	50.2	7.7	37.9	1.0	0.2	3.0
1504	955	47.5	3.6	42.6	0.0	5.1	1.2
1505	244	80.3	1.6	13.5	0.4	2.9	1.2
1506	196	84.7	1.0	12.8	0.0	0.0	1.5
1507	230	95.2	0.9	2.2	0.0	0.4	1.3
1508	87	*	*	*	*	*	*
1509	236	51.3	3.4	45.3	0.0	0.0	0.0
1510	72	*	*	*	*	*	*
1511	195	93.8	0.5	3.6	0.0	0.0	2.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 4  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
1512	126	93.7	0.8	4.0	0.0	0.8	0.8
1701	248	63.7	1.2	1.2	19.0	0.4	14.5
1712	44	*	*	*	*	*	*
1950	253	95.7	0.8	2.8	0.4	0.0	0.4
1952	60	*	*	*	*	*	*
1953	139	97.8	0.0	2.2	0.0	0.0	0.0
1954	146	89.7	1.4	8.9	0.0	0.0	0.0
1955	98	*	*	*	*	*	*
1957	135	96.3	0.0	3.7	0.0	0.0	0.0
1959	66	*	*	*	*	*	*
1963	69	*	*	*	*	*	*
2150	254	92.9	0.4	0.0	0.0	0.8	5.9
2151	116	94.0	0.0	5.2	0.0	0.0	0.9
2251	188	95.7	0.0	0.5	1.1	2.1	0.5
2252	67	*	*	*	*	*	*
2551	1,343	31.6	66.0	1.9	0.1	0.1	0.2
2552	336	27.1	67.6	4.5	0.0	0.3	0.6
2554	45	*	*	*	*	*	*
2555	1,533	79.6	17.7	1.9	0.1	0.5	0.1
3001	409	15.2	70.9	12.7	0.0	0.7	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 5  
Run Date: 10 / 11 / 07

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan Native %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other/Unknown %</b>	
3002	89	*	*	*	*	*	*	
3003	7	*	*	*	*	*	*	
3004	89	*	*	*	*	*	*	
3007	12	*	*	*	*	*	*	
3009	3	*	*	*	*	*	*	
3201	240	94.6	0.0	5.0	0.0	0.0	0.4	
3202	1	*	*	*	*	*	*	
3301	1,415	54.5	26.9	14.2	0.4	3.1	0.9	
3302	144	27.1	41.0	20.8	0.7	9.0	1.4	
3303	22	*	*	*	*	*	*	
3304	56	*	*	*	*	*	*	
3305	31	*	*	*	*	*	*	
3306	52	*	*	*	*	*	*	
3308	17	*	*	*	*	*	*	
3309	15	*	*	*	*	*	*	
3311	7	*	*	*	*	*	*	
3550	30	*	*	*	*	*	*	
3551	10	*	*	*	*	*	*	
3552	14	*	*	*	*	*	*	
3650	7	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 6  
Run Date: 10 / 11 / 07

<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>					
		<b>White, Not Hispanic</b> <b>%</b>	<b>Black, Not Hispanic</b> <b>%</b>	<b>Hispanic</b> <b>%</b>	<b>American Indian/Alaskan Native</b> <b>%</b>	<b>Asian/Pacific Islander</b> <b>%</b>	<b>All Other/Unknown</b> <b>%</b>
3701	271	89.7	1.1	6.6	0.7	0.7	1.1
3702	88	*	*	*	*	*	*
3703	36	*	*	*	*	*	*
3704	40	*	*	*	*	*	*
3805	1,121	77.9	15.3	5.6	0.5	0.1	0.6
3950	707	66.9	22.2	7.6	0.0	1.8	1.4
3952	22	*	*	*	*	*	*
3953	31	*	*	*	*	*	*
3954	77	*	*	*	*	*	*
4001	355	23.7	62.0	12.4	0.0	0.8	1.1
4002	321	52.0	32.1	12.8	0.0	1.2	1.9
4103	12	*	*	*	*	*	*
4106	578	6.9	49.7	41.5	0.0	1.2	0.7
4107	1,108	55.4	29.1	11.6	0.4	1.5	2.0
4109	399	94.2	1.0	2.3	0.3	0.5	1.8
4110	652	36.2	8.3	52.6	0.6	0.5	1.8
4111	17	*	*	*	*	*	*
4112	286	1.0	0.3	98.3	0.0	0.0	0.3
4113	320	7.8	16.9	73.4	0.0	0.9	0.9
4120	734	65.5	9.5	20.3	0.4	3.0	1.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 7  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
4160	783	45.3	32.6	12.3	0.4	7.5	1.9
4351	259	93.4	1.2	3.5	0.4	0.8	0.8
4352	75	*	*	*	*	*	*
4353	187	73.8	0.5	23.0	0.5	0.0	2.1
4354	308	89.6	5.8	2.6	0.0	0.6	1.3
4355	131	93.1	0.0	0.8	0.8	0.0	5.3
4356	219	86.8	1.4	10.5	0.0	0.5	0.9
4357	2	*	*	*	*	*	*
4358	88	*	*	*	*	*	*
4359	137	69.3	0.0	29.9	0.0	0.7	0.0
4360	79	*	*	*	*	*	*
4361	104	95.2	3.8	0.0	0.0	0.0	1.0
4362	133	98.5	0.0	0.8	0.0	0.0	0.8
4363	336	94.3	0.6	2.1	0.3	0.6	2.1
4401	445	91.2	0.4	5.4	0.4	0.4	2.0
4501	345	96.5	0.6	2.6	0.0	0.3	0.0
4601	556	78.1	2.0	19.1	0.0	0.2	0.7
4701	390	95.1	0.0	3.3	0.0	1.5	0.0
5050	411	76.6	18.5	1.9	0.2	1.5	1.2
5051	1,057	71.8	15.3	8.5	0.0	2.9	1.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 8  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %	
5052	632	73.7	16.8	2.7	0.2	5.5	1.1	
5054	87	*	*	*	*	*	*	
5202	125	92.8	0.8	0.0	0.8	0.8	4.8	
5203	61	*	*	*	*	*	*	
5204	56	*	*	*	*	*	*	
5205	81	*	*	*	*	*	*	
5601	379	97.6	1.1	0.5	0.3	0.0	0.5	
5801	779	89.1	5.0	4.5	0.1	0.5	0.8	
5803	3	*	*	*	*	*	*	
6103	1,585	61.5	27.9	6.9	0.7	0.7	2.4	
6301	1,719	45.1	28.5	23.5	0.0	2.2	0.7	
6312	1,029	41.3	53.7	1.2	0.1	2.5	1.2	
6316	403	79.4	9.9	7.9	0.0	0.5	2.2	
6320	93	*	*	*	*	*	*	
6325	54	*	*	*	*	*	*	
6330	109	31.2	51.4	13.8	0.0	2.8	0.9	
6335	209	81.3	12.0	2.9	1.0	2.9	0.0	
6340	168	50.0	12.5	33.3	0.0	3.6	0.6	
7304	100	94.0	1.0	4.0	0.0	0.0	1.0	
7306	991	49.5	34.9	14.2	0.0	0.6	0.7	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 9  
Run Date: 10 / 11 / 07

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan Native %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other/Unknown %</b>	
7308	70	*	*	*	*	*	*	
7401	797	83.9	6.6	4.9	0.6	0.4	3.5	
7402	116	88.8	1.7	4.3	0.0	0.0	5.2	
7403	37	*	*	*	*	*	*	
7404	48	*	*	*	*	*	*	
7601	234	95.3	0.4	3.0	0.0	0.0	1.3	
7850	436	94.7	0.5	3.0	0.0	0.2	1.6	
7851	58	*	*	*	*	*	*	
7852	33	*	*	*	*	*	*	
8150	37	*	*	*	*	*	*	
8151	918	38.9	44.4	13.9	0.2	1.7	0.8	
8152	214	41.6	45.8	7.5	0.0	1.4	3.7	
8153	9	*	*	*	*	*	*	
8154	134	33.6	20.9	7.5	0.0	37.3	0.7	
8155	32	*	*	*	*	*	*	
8156	12	*	*	*	*	*	*	
8157	3	*	*	*	*	*	*	
8201	868	62.3	28.6	3.8	0.5	2.0	2.9	
8202	1,167	85.9	6.0	5.1	0.3	2.1	0.6	
8253	2	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 10  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
8254	4	*	*	*	*	*	*
8255	1,139	76.5	20.8	1.8	0.0	0.6	0.3
8257	242	48.3	25.2	0.0	0.0	25.6	0.8
8258	564	72.2	22.0	3.0	0.2	1.1	1.6
8901	387	0.8	96.9	0.8	0.0	1.0	0.5
8902	168	3.0	97.0	0.0	0.0	0.0	0.0
8903	463	4.1	93.3	0.9	0.0	0.6	1.1
8904	284	7.0	67.3	1.1	1.4	22.5	0.7
8905	786	20.4	31.9	34.5	0.6	10.3	2.3
9004	258	0.4	98.8	0.4	0.0	0.4	0.0
9010	78	*	*	*	*	*	*
9023	37	*	*	*	*	*	*
9040	176	1.1	98.3	0.0	0.0	0.0	0.6
9045	422	9.5	55.5	34.6	0.0	0.5	0.0
9050	771	13.7	49.3	35.7	0.0	1.3	0.0
9051	1,043	6.8	7.6	84.9	0.2	0.0	0.6
9053	1,018	0.6	98.6	0.4	0.0	0.4	0.0
9054	183	2.2	91.8	2.7	0.5	1.6	1.1
9058	188	79.3	17.6	0.5	0.0	2.7	0.0
9066	916	1.0	98.7	0.2	0.0	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**

Page: 11  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
9068	612	2.8	96.2	0.0	0.0	1.0	0.0
9069	807	1.7	98.0	0.0	0.0	0.1	0.1
9070	799	2.4	87.6	0.1	0.5	8.9	0.5
9072	580	2.2	92.9	0.2	0.0	4.7	0.0
9074	527	1.1	92.0	1.3	0.0	0.0	5.5
9079	475	0.2	98.7	0.0	0.0	0.6	0.4
9080	775	27.4	68.5	3.6	0.0	0.0	0.5
9151	1	*	*	*	*	*	*
Unknown Clinics	375	18.9	65.3	6.7	0.0	0.8	8.3
<b>Michigan</b>	<b>61,323</b>	<b>56.0</b>	<b>29.0</b>	<b>11.6</b>	<b>0.4</b>	<b>1.7</b>	<b>1.3</b>
<b>Nation (Prior Year)</b>	<b>857,244</b>	<b>48.9</b>	<b>23.5</b>	<b>20.0</b>	<b>1.3</b>	<b>2.4</b>	<b>3.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 12  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0101	58	*	*	*	*	*	*	*	*
0102	159	0.0	8.2	10.7	66.0	13.8	1.3	0.0	
0103	168	0.0	7.7	13.7	67.3	10.1	1.2	0.0	
0104	58	*	*	*	*	*	*	*	*
0250	64	*	*	*	*	*	*	*	*
0251	41	*	*	*	*	*	*	*	*
0252	55	*	*	*	*	*	*	*	*
0253	52	*	*	*	*	*	*	*	*
0401	195	0.0	7.7	9.7	68.2	12.8	1.5	0.0	
0402	172	0.0	5.8	10.5	64.0	19.8	0.0	0.0	
0403	60	*	*	*	*	*	*	*	*
0404	55	*	*	*	*	*	*	*	*
0501	203	0.5	6.9	14.3	64.5	13.3	0.5	0.0	
0502	2	*	*	*	*	*	*	*	*
0503	68	*	*	*	*	*	*	*	*
0504	239	0.0	4.2	11.7	66.5	16.7	0.8	0.0	
0505	155	0.0	5.2	12.9	71.0	9.7	1.3	0.0	
0506	74	*	*	*	*	*	*	*	*
0650	382	0.0	3.9	13.6	68.3	13.4	0.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 13  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0651	252	0.8	7.1	17.9	61.1	12.7	0.4	0.0	
0652	136	0.0	5.1	15.4	62.5	16.2	0.7	0.0	
0653	147	0.7	6.8	19.7	57.8	14.3	0.7	0.0	
0654	128	0.8	7.0	14.8	64.1	13.3	0.0	0.0	
0655	130	0.8	6.2	13.8	65.4	12.3	1.5	0.0	
0657	42	*	*	*	*	*	*	*	*
0701	173	0.0	4.6	9.8	63.6	19.7	2.3	0.0	
0702	38	*	*	*	*	*	*	*	*
0703	94	*	*	*	*	*	*	*	*
0704	32	*	*	*	*	*	*	*	*
0705	59	*	*	*	*	*	*	*	*
0850	399	0.0	4.5	12.5	68.7	13.8	0.5	0.0	
0851	337	0.3	8.6	18.4	58.5	12.8	1.5	0.0	
0901	675	0.1	7.6	11.6	63.0	16.0	1.8	0.0	
0902	21	*	*	*	*	*	*	*	*
1051	540	0.4	5.7	10.6	61.3	21.3	0.7	0.0	
1101	724	0.7	9.8	13.7	60.2	14.9	0.7	0.0	
1102	336	0.0	8.9	14.0	62.2	14.3	0.6	0.0	
1106	49	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 14  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1250	387	0.3	9.0	15.8	63.3	10.9	0.8	0.0	
1251	283	1.1	7.8	14.5	62.2	13.1	1.4	0.0	
1252	293	0.0	5.1	11.3	68.6	14.7	0.3	0.0	
1253	278	0.4	6.8	16.9	61.9	13.3	0.7	0.0	
1350	905	0.2	10.2	16.5	59.9	12.5	0.8	0.0	
1351	56	*	*	*	*	*	*	*	*
1352	153	0.0	9.8	16.3	59.5	13.1	1.3	0.0	
1401	110	0.0	5.5	10.9	69.1	12.7	1.8	0.0	
1402	61	*	*	*	*	*	*	*	*
1403	8	*	*	*	*	*	*	*	*
1404	8	*	*	*	*	*	*	*	*
1501	423	0.0	6.9	11.6	65.0	16.1	0.5	0.0	
1502	45	*	*	*	*	*	*	*	*
1503	404	0.0	9.2	14.1	56.2	19.8	0.7	0.0	
1504	955	0.7	7.7	15.5	59.2	16.2	0.6	0.0	
1505	244	0.4	4.5	10.7	66.8	17.6	0.0	0.0	
1506	196	0.0	5.1	13.3	64.8	15.8	1.0	0.0	
1507	230	0.0	3.9	10.9	68.7	15.2	1.3	0.0	
1508	87	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 15  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1509	236	0.4	6.4	12.7	65.3	14.4	0.8	0.0	
1510	72	*	*	*	*	*	*	*	*
1511	195	0.5	6.2	15.4	68.7	9.2	0.0	0.0	
1512	126	0.0	3.2	12.7	62.7	19.8	1.6	0.0	
1701	248	0.0	5.6	11.3	63.3	18.5	1.2	0.0	
1712	44	*	*	*	*	*	*	*	*
1950	253	0.4	7.1	16.6	65.6	9.5	0.8	0.0	
1952	60	*	*	*	*	*	*	*	*
1953	139	0.7	3.6	12.2	74.1	8.6	0.7	0.0	
1954	146	1.4	4.8	13.0	74.7	6.2	0.0	0.0	
1955	98	*	*	*	*	*	*	*	*
1957	135	0.0	3.0	15.6	68.9	12.6	0.0	0.0	
1959	66	*	*	*	*	*	*	*	*
1963	69	*	*	*	*	*	*	*	*
2150	254	0.0	5.9	12.2	67.7	13.0	1.2	0.0	
2151	116	0.0	5.2	16.4	64.7	13.8	0.0	0.0	
2251	188	0.0	5.9	10.1	64.9	18.6	0.5	0.0	
2252	67	*	*	*	*	*	*	*	*
2551	1,343	0.8	10.1	14.7	57.2	16.5	0.7	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 16  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
2552	336	1.5	9.2	17.0	58.0	13.7	0.6	0.0	
2554	45	*	*	*	*	*	*	*	*
2555	1,533	0.4	5.4	11.6	66.2	15.7	0.7	0.0	
3001	409	1.2	14.9	14.7	53.1	15.9	0.2	0.0	
3002	89	*	*	*	*	*	*	*	*
3003	7	*	*	*	*	*	*	*	*
3004	89	*	*	*	*	*	*	*	*
3007	12	*	*	*	*	*	*	*	*
3009	3	*	*	*	*	*	*	*	*
3201	240	0.0	4.6	7.9	62.1	23.3	2.1	0.0	
3202	1	*	*	*	*	*	*	*	*
3301	1,415	0.1	6.5	14.6	63.7	14.4	0.7	0.0	
3302	144	0.7	5.6	18.1	56.9	18.1	0.7	0.0	
3303	22	*	*	*	*	*	*	*	*
3304	56	*	*	*	*	*	*	*	*
3305	31	*	*	*	*	*	*	*	*
3306	52	*	*	*	*	*	*	*	*
3308	17	*	*	*	*	*	*	*	*
3309	15	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 17  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
3311	7	*	*	*	*	*	*	*	*
3550	30	*	*	*	*	*	*	*	*
3551	10	*	*	*	*	*	*	*	*
3552	14	*	*	*	*	*	*	*	*
3650	7	*	*	*	*	*	*	*	*
3701	271	0.0	8.1	17.3	61.3	12.2	1.1	0.0	
3702	88	*	*	*	*	*	*	*	*
3703	36	*	*	*	*	*	*	*	*
3704	40	*	*	*	*	*	*	*	*
3805	1,121	0.1	10.4	16.1	60.3	12.7	0.4	0.0	
3950	707	0.1	6.9	11.7	64.4	15.4	1.4	0.0	
3952	22	*	*	*	*	*	*	*	*
3953	31	*	*	*	*	*	*	*	*
3954	77	*	*	*	*	*	*	*	*
4001	355	0.3	9.9	14.9	63.4	11.5	0.0	0.0	
4002	321	0.3	8.1	17.1	61.1	12.5	0.9	0.0	
4103	12	*	*	*	*	*	*	*	*
4106	578	0.7	9.0	15.1	57.4	16.4	1.4	0.0	
4107	1,108	0.5	6.5	14.1	60.5	17.3	1.2	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 18  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4109	399	0.3	6.5	12.8	60.9	18.0	1.5	0.0	
4110	652	0.8	6.3	12.1	63.2	17.2	0.5	0.0	
4111	17	*	*	*	*	*	*	*	*
4112	286	0.0	3.1	9.4	54.9	29.7	2.8	0.0	
4113	320	1.9	10.0	13.1	56.9	17.2	0.9	0.0	
4120	734	0.3	7.6	13.2	60.6	16.2	2.0	0.0	
4160	783	0.3	6.8	12.1	62.3	17.2	1.3	0.0	
4351	259	0.4	5.4	20.8	62.5	10.0	0.8	0.0	
4352	75	*	*	*	*	*	*	*	*
4353	187	1.1	11.8	17.6	59.9	9.1	0.5	0.0	
4354	308	0.0	8.1	18.5	60.1	12.7	0.6	0.0	
4355	131	0.8	6.1	12.2	67.2	13.0	0.8	0.0	
4356	219	0.0	5.0	12.8	64.4	16.4	1.4	0.0	
4357	2	*	*	*	*	*	*	*	*
4358	88	*	*	*	*	*	*	*	*
4359	137	0.0	9.5	14.6	59.1	15.3	1.5	0.0	
4360	79	*	*	*	*	*	*	*	*
4361	104	1.0	11.5	10.6	64.4	12.5	0.0	0.0	
4362	133	0.0	4.5	12.8	73.7	8.3	0.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 19  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4363	336	0.0	6.0	17.0	67.9	8.3	0.9	0.0	
4401	445	0.0	4.5	16.2	59.1	19.1	1.1	0.0	
4501	345	0.0	7.5	11.3	66.7	14.5	0.0	0.0	
4601	556	0.2	7.7	13.7	64.7	13.1	0.5	0.0	
4701	390	0.0	4.4	18.7	61.0	14.6	1.3	0.0	
5050	411	0.2	4.1	10.7	58.9	24.1	1.9	0.0	
5051	1,057	0.1	5.1	11.7	62.6	18.4	2.0	0.0	
5052	632	0.2	4.3	11.7	58.2	23.9	1.7	0.0	
5054	87	*	*	*	*	*	*	*	
5202	125	0.0	2.4	12.8	69.6	14.4	0.8	0.0	
5203	61	*	*	*	*	*	*	*	
5204	56	*	*	*	*	*	*	*	
5205	81	*	*	*	*	*	*	*	
5601	379	0.3	6.6	15.0	63.3	13.7	1.1	0.0	
5801	779	0.3	5.1	15.4	64.2	14.4	0.6	0.0	
5803	3	*	*	*	*	*	*	*	
6103	1,585	0.3	8.6	14.6	62.5	13.7	0.4	0.0	
6301	1,719	0.3	6.7	13.0	60.7	17.9	1.4	0.0	
6312	1,029	0.1	4.5	9.5	56.2	27.6	2.1	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 20  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
6316	403	0.0	3.2	12.4	67.2	15.9	1.2	0.0	
6320	93	*	*	*	*	*	*	*	*
6325	54	*	*	*	*	*	*	*	*
6330	109	0.0	7.3	16.5	58.7	12.8	4.6	0.0	
6335	209	0.0	3.8	10.5	60.3	21.1	4.3	0.0	
6340	168	0.6	1.8	14.9	64.3	18.5	0.0	0.0	
7304	100	0.0	6.0	10.0	65.0	17.0	2.0	0.0	
7306	991	0.0	7.5	14.9	60.3	16.1	1.1	0.0	
7308	70	*	*	*	*	*	*	*	*
7401	797	0.3	6.3	13.6	65.1	13.8	1.0	0.0	
7402	116	0.0	6.0	13.8	62.1	17.2	0.9	0.0	
7403	37	*	*	*	*	*	*	*	*
7404	48	*	*	*	*	*	*	*	*
7601	234	0.0	8.1	12.4	63.7	15.8	0.0	0.0	
7850	436	0.0	4.8	15.8	67.4	10.6	1.4	0.0	
7851	58	*	*	*	*	*	*	*	*
7852	33	*	*	*	*	*	*	*	*
8150	37	*	*	*	*	*	*	*	*
8151	918	0.1	2.1	7.4	66.9	23.1	0.4	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 21  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
8152	214	1.9	26.6	43.5	28.0	0.0	0.0	0.0	
8153	9	*	*	*	*	*	*	*	*
8154	134	0.0	3.0	3.0	56.7	36.6	0.7	0.0	
8155	32	*	*	*	*	*	*	*	*
8156	12	*	*	*	*	*	*	*	*
8157	3	*	*	*	*	*	*	*	*
8201	868	0.5	6.7	13.8	57.7	20.4	0.9	0.0	
8202	1,167	0.1	2.7	9.2	61.0	24.4	2.7	0.0	
8253	2	*	*	*	*	*	*	*	*
8254	4	*	*	*	*	*	*	*	*
8255	1,139	0.1	6.9	12.3	62.3	17.3	1.1	0.0	
8257	242	0.0	3.7	9.5	60.3	21.9	4.5	0.0	
8258	564	0.2	9.9	12.9	56.7	18.8	1.4	0.0	
8901	387	1.3	13.4	13.4	56.6	14.7	0.5	0.0	
8902	168	1.2	4.8	11.9	57.1	21.4	3.6	0.0	
8903	463	0.4	11.4	17.1	46.4	23.1	1.5	0.0	
8904	284	2.5	6.7	10.2	61.3	18.0	1.4	0.0	
8905	786	0.6	7.9	12.7	51.1	25.7	1.9	0.0	
9004	258	1.9	10.1	12.0	56.2	18.2	1.6	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 22  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
9010	78	*	*	*	*	*	*	*	*
9023	37	*	*	*	*	*	*	*	*
9040	176	2.8	10.8	15.9	52.3	16.5	1.7	0.0	
9045	422	3.6	17.1	13.0	46.9	18.2	1.2	0.0	
9050	771	0.8	8.8	12.1	54.0	22.2	2.2	0.0	
9051	1,043	0.5	7.6	10.7	58.9	21.4	1.0	0.0	
9053	1,018	0.5	8.5	15.4	53.9	20.1	1.5	0.0	
9054	183	1.1	4.9	11.5	61.7	20.2	0.5	0.0	
9058	188	0.0	3.7	8.0	55.9	29.3	3.2	0.0	
9066	916	0.4	10.5	15.5	50.8	20.9	1.9	0.1	
9068	612	0.5	7.4	13.2	60.8	16.5	1.6	0.0	
9069	807	0.9	8.1	13.0	58.6	18.3	1.0	0.1	
9070	799	0.5	9.8	14.0	56.4	18.3	1.0	0.0	
9072	580	1.6	10.7	15.2	52.6	19.1	0.9	0.0	
9074	527	0.6	7.0	13.1	58.8	18.4	2.1	0.0	
9079	475	0.6	5.3	9.9	56.4	25.9	1.9	0.0	
9080	775	0.4	7.0	12.6	55.1	23.6	1.3	0.0	
9151	1	*	*	*	*	*	*	*	*
Unknown Clinics	375	1.3	10.1	13.3	52.0	21.3	1.9	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Page: 23  
Run Date: 10 / 11 / 07

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
<b>Michigan</b>	<b>61,323</b>	<b>0.4</b>	<b>7.2</b>	<b>13.4</b>	<b>60.7</b>	<b>17.2</b>	<b>1.2</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>857,244</b>	<b>0.4</b>	<b>6.6</b>	<b>13.0</b>	<b>60.4</b>	<b>18.2</b>	<b>1.3</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 24  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
0101	57	* * * *	*	44	* * * *	*	7	* *	16	* *		
0102	153	7.2 (9)	50.3 (100)	129	26.4 (62)	53.5 (114)	24	* *	31	* *		
0103	163	14.7 (115)	49.7 (94)	139	25.9 (55)	47.5 (78)	17	* *	31	* *		
0104	57	* * * *	*	50	* * * *	*	5	* *	15	* *		
0250	63	* * * *	*	50	* * * *	*	6	* *	19	* *		
0251	39	* * * *	*	36	* * * *	*	5	* *	9	* *		
0252	52	* * * *	*	40	* * * *	*	6	* *	16	* *		
0253	49	* * * *	*	43	* * * *	*	4	* *	9	* *		
0401	192	12.5 (92)	48.4 (84)	168	24.4 (47)	50.0 (101)	22	* *	84	* *		
0402	170	13.5 (106)	46.5 (63)	144	26.4 (62)	41.0 (32)	25	* *	36	* *		
0403	59	* * * *	*	55	* * * *	*	9	* *	17	* *		
0404	54	* * * *	*	51	* * * *	*	9	* *	18	* *		
0501	196	14.3 (111)	37.2 (7)	182	17.6 (6)	48.4 (88)	31	* *	64	* *		
0502	2	* * * *	*	2	* * * *	*	1	* *	0	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 25  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
0503	58	* * * *		55	* * * *		13	* *	21	* *		
0504	236	12.7 (93)	35.2 (5)	218	33.0 (91)	38.5 (22)	25	* *	56	* *		
0505	153	10.5 (53)	47.7 (76)	140	62.1 (119)	20.0 (2)	18	* *	45	* *		
0506	68	* * * *		59	* * * *		7	* *	15	* *		
0650	366	10.1 (40)	47.8 (77)	333	26.1 (58)	46.8 (72)	51	* *	103	14.6 (3)		
0651	242	10.7 (60)	49.2 (90)	209	24.4 (47)	48.8 (91)	28	* *	85	* *		
0652	128	10.2 (42)	44.5 (42)	106	32.1 (88)	43.4 (48)	16	* *	28	* *		
0653	144	10.4 (50)	47.2 (70)	130	33.1 (93)	38.5 (22)	30	* *	19	* *		
0654	125	14.4(113)	48.8 (87)	94	* * * *		14	* *	41	* *		
0655	126	9.5 (31)	47.6 (74)	111	26.1 (58)	42.3 (43)	22	* *	44	* *		
0657	41	* * * *		36	* * * *		5	* *	19	* *		
0701	166	21.1(123)	30.1 (3)	140	23.6 (38)	44.3 (57)	25	* *	55	* *		
0702	37	* * * *		33	* * * *		6	* *	7	* *		
0703	86	* * * *		79	* * * *		8	* *	20	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 26  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
0704	31	* * * *		28	* * * *		4	* *	7	* *
0705	55	* * * *		49	* * * *		8	* *	18	* *
0850	382	10.5 (53)	47.6 (74)	330	33.0 (91)	43.6 (50)	58	* *	117	35.9 (28)
0851	326	12.0 (84)	42.9 (30)	290	38.3 (103)	37.6 (21)	33	* *	85	* *
0901	642	14.5(114)	46.6 (64)	578	26.0 (57)	47.1 (77)	107	15.0 (1)	189	10.6 (1)
0902	21	* * * *		17	* * * *		5	* *	5	* *
1051	527	10.2 (42)	42.7 (28)	455	20.0 (15)	47.9 (83)	71	* *	183	27.9 (17)
1101	697	11.6 (79)	50.2 (99)	581	25.3 (52)	50.4 (103)	155	48.4 (30)	187	44.9 (39)
1102	314	12.1 (87)	45.2 (52)	268	23.5 (36)	51.5 (108)	58	* *	108	47.2 (41)
1106	45	* * * *		42	* * * *		13	* *	20	* *
1250	373	13.1(101)	42.1 (26)	330	30.0 (74)	47.6 (79)	58	* *	88	* *
1251	273	12.8 (96)	46.9 (67)	246	30.1 (75)	45.5 (65)	48	* *	85	* *
1252	283	11.3 (74)	52.7 (115)	237	28.7 (71)	44.7 (59)	48	* *	66	* *
1253	269	10.8 (63)	45.4 (55)	227	40.1 (106)	38.8 (24)	39	* *	63	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 27  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
1350	874	10.2 (42)	46.6 (64)	760	23.4 (35)	48.7 (90)	162	29.6 (13)	140	56.4 (46)
1351	54	* * * *		47	* * * *		10	* *	10	* *
1352	148	8.1 (15)	50.0 (97)	128	24.2 (44)	46.1 (67)	26	* *	26	* *
1401	106	13.2(103)	48.1 (81)	90	* * * *		12	* *	30	* *
1402	59	* * * *		52	* * * *		9	* *	11	* *
1403	8	* * * *		7	* * * *		1	* *	2	* *
1404	6	* * * *		2	* * * *		2	* *	1	* *
1501	394	9.9 (36)	55.3 (120)	333	30.6 (80)	35.7 (17)	48	* *	75	* *
1502	45	* * * *		36	* * * *		9	* *	8	* *
1503	360	9.4 (29)	51.1 (104)	289	56.7 (117)	26.6 (6)	58	* *	60	* *
1504	917	11.1 (68)	41.2 (21)	780	44.1 (110)	33.1 (10)	165	30.3 (15)	214	37.4 (32)
1505	234	9.8 (34)	38.9 (10)	207	19.8 (13)	46.9 (74)	41	* *	36	* *
1506	183	12.0 (84)	41.0 (17)	160	22.5 (28)	50.6 (105)	25	* *	24	* *
1507	220	12.3 (90)	45.0 (49)	196	12.2 (1)	49.0 (94)	33	* *	41	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 28  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
1508	52	* * * *	*	34	* * * *	*	8	* *	11	* *		
1509	224	3.1 (1)	58.5 (125)	177	38.4 (104)	39.0 (26)	23	* *	26	* *		
1510	68	* * * *	*	52	* * * *	*	11	* *	14	* *		
1511	190	10.0 (38)	52.6 (114)	164	17.7 (7)	52.4 (112)	35	* *	26	* *		
1512	121	18.2 (122)	46.3 (60)	107	22.4 (27)	43.9 (51)	28	* *	23	* *		
1701	246	8.9 (25)	50.4 (101)	219	19.2 (8)	50.7 (106)	26	* *	39	* *		
1712	43	* * * *	*	39	* * * *	*	7	* *	4	* *		
1950	251	15.5(116)	44.6 (43)	217	16.6 (3)	53.5 (114)	40	* *	61	* *		
1952	60	* * * *	*	54	* * * *	*	11	* *	15	* *		
1953	138	16.7(120)	42.8 (29)	127	20.5 (17)	50.4 (103)	18	* *	32	* *		
1954	140	8.6 (21)	57.1 (124)	113	31.0 (82)	44.2 (55)	12	* *	27	* *		
1955	96	* * * *	*	80	* * * *	*	12	* *	17	* *		
1957	128	11.7 (80)	40.6 (14)	116	19.8 (13)	49.1 (96)	24	* *	29	* *		
1959	64	* * * *	*	54	* * * *	*	15	* *	14	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 29  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
1963	67	* * * *		55	* * * *		9	* *	12	* *		
2150	248	10.5 (53)	46.0 (59)	210	21.9 (23)	54.8 (117)	32	* *	57	* *		
2151	108	8.3 (17)	43.5 (34)	91	* * * *		14	* *	23	* *		
2251	172	10.5 (53)	45.3 (54)	153	23.5 (36)	48.4 (88)	22	* *	57	* *		
2252	64	* * * *		51	* * * *		1	* *	12	* *		
2551	1,311	11.2 (71)	48.9 (88)	1,120	24.2 (44)	49.6 (97)	274	49.6 (31)	294	41.2 (37)		
2552	328	8.8 (23)	47.3 (71)	275	19.3 (10)	49.8 (100)	63	* *	47	* *		
2554	45	* * * *		39	* * * *		2	* *	6	* *		
2555	1,480	13.1 (101)	46.3 (60)	1,246	23.2 (34)	49.0 (94)	281	35.6 (22)	334	28.7 (18)		
3001	387	9.0 (26)	51.2 (105)	324	30.2 (77)	42.6 (44)	60	* *	192	31.8 (23)		
3002	83	* * * *		75	* * * *		11	* *	47	* *		
3003	6	* * * *		4	* * * *		1	* *	1	* *		
3004	83	* * * *		71	* * * *		6	* *	14	* *		
3007	11	* * * *		8	* * * *		1	* *	6	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 30  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
3009	2	*	*	1	*	*	0	*	0	*	*	
3201	234	10.3 (48)	50.0 (97)	202	23.8 (42)	44.1 (53)	34	*	32	*	*	
3202	0	*	*	*	*	*	0	*	0	*	*	
3301	1,361	10.9 (65)	43.3 (32)	1,175	45.2 (111)	34.8 (15)	287	28.2 (9)	422	45.5 (40)		
3302	139	5.0 (3)	56.8 (123)	120	32.5 (90)	44.2 (55)	23	*	51	*	*	
3303	20	*	*	*	*	*	3	*	7	*	*	
3304	55	*	*	*	*	*	7	*	12	*	*	
3305	30	*	*	*	*	*	3	*	6	*	*	
3306	48	*	*	*	*	*	10	*	17	*	*	
3308	15	*	*	*	*	*	3	*	6	*	*	
3309	14	*	*	*	*	*	1	*	4	*	*	
3311	6	*	*	*	*	*	0	*	2	*	*	
3550	29	*	*	*	*	*	1	*	4	*	*	
3551	9	*	*	*	*	*	2	*	4	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 31  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
3552	14	* * * *	*	11	* * * *	*	3	* *	5	* *		
3650	7	* * * *	*	1	* * * *	*	3	* *	0	* *		
3701	266	9.4 (29)	44.7 (44)	228	26.3 (60)	46.5 (69)	41	* *	52	* *		
3702	86	* * * *	*	71	* * * *	*	20	* *	13	* *		
3703	36	* * * *	*	34	* * * *	*	7	* *	8	* *		
3704	38	* * * *	*	29	* * * *	*	8	* *	3	* *		
3805	1,088	11.1 (68)	44.8 (47)	975	27.7 (67)	47.7 (82)	157	28.7 (10)	446	40.4 (36)		
3950	697	9.8 (34)	49.8 (96)	602	20.3 (16)	53.2 (113)	99	* *	148	39.2 (35)		
3952	22	* * * *	*	19	* * * *	*	2	* *	4	* *		
3953	31	* * * *	*	28	* * * *	*	6	* *	7	* *		
3954	76	* * * *	*	64	* * * *	*	11	* *	13	* *		
4001	341	8.8 (23)	52.5 (113)	258	31.8 (87)	45.0 (60)	42	* *	47	* *		
4002	308	11.4 (75)	45.5 (56)	247	27.9 (68)	42.1 (39)	42	* *	47	* *		
4103	11	* * * *	*	9	* * * *	*	2	* *	2	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 32  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
4106	562	6.8 (5)	53.0 (116)	488	26.4 (62)	40.8 (30)	97	* *	121	36.4 (30)
4107	1,049	11.2 (71)	45.1 (51)	906	25.9 (55)	45.0 (60)	190	24.2 (5)	276	26.1 (15)
4109	381	12.9 (98)	41.2 (21)	343	23.0 (31)	46.6 (71)	65	* *	115	16.5 (4)
4110	628	8.3 (17)	48.1 (81)	532	26.3 (60)	41.2 (33)	63	* *	146	36.3 (29)
4111	15	* * * *		12	* * * *		3	* *	5	* *
4112	250	3.2 (2)	51.6 (107)	226	59.7 (118)	15.0 (1)	44	* *	112	60.7 (47)
4113	320	7.2 (9)	44.7 (44)	285	31.2 (84)	31.9 (9)	31	* *	73	* *
4120	718	12.0 (84)	45.7 (58)	636	22.0 (24)	51.9 (110)	122	24.6 (6)	181	22.7 (10)
4160	762	11.4 (75)	41.5 (24)	663	25.8 (54)	43.4 (48)	124	29.8 (14)	203	19.2 (8)
4351	254	11.8 (82)	42.5 (27)	228	23.7 (41)	49.6 (97)	29	* *	66	* *
4352	71	* * * *		60	* * * *		11	* *	20	* *
4353	182	10.4 (50)	51.6 (107)	148	19.6 (12)	56.8 (119)	23	* *	42	* *
4354	300	10.0 (38)	49.0 (89)	254	17.3 (5)	48.8 (91)	47	* *	79	* *
4355	122	13.9 (109)	41.0 (17)	110	23.6 (38)	53.6 (116)	15	* *	33	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 33  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)													
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)				
4356	210	11.0 (67)	44.3 (37)	193	22.3 (26)	52.3 (111)	37	* *	60	* *				
4357	0	*	*	*	*	*	0	*	*	0	*	*		
4358	75	*	*	*	*	*	10	*	*	21	*	*		
4359	135	13.3 (104)	40.7 (15)	126	29.4 (73)	42.1 (39)	11	*	*	26	*	*		
4360	78	*	*	*	*	*	8	*	*	25	*	*		
4361	102	12.7 (93)	39.2 (13)	84	*	*	*	*	15	*	*	30	*	*
4362	131	9.9 (36)	44.3 (37)	125	19.2 (8)	41.6 (35)	18	*	*	41	*	*		
4363	333	10.5 (53)	44.7 (44)	296	19.3 (10)	50.0 (101)	41	*	*	102	23.5 (11)			
4401	219	16.4 (119)	44.3 (37)	196	28.1 (70)	43.9 (51)	69	*	*	177	37.9 (33)			
4501	308	14.3 (111)	47.4 (72)	275	25.5 (53)	43.3 (47)	45	*	*	80	*	*		
4601	551	12.2 (89)	46.3 (60)	480	22.7 (29)	46.9 (74)	44	*	*	99	*	*		
4701	386	11.7 (80)	39.1 (11)	337	21.1 (21)	55.2 (118)	90	*	*	137	13.1 (2)			
5050	403	10.7 (60)	48.4 (84)	88	*	*	*	*	101	28.7 (10)	64	*	*	
5051	1,014	14.0 (110)	41.9 (25)	321	30.8 (81)	41.7 (36)	257	32.3 (17)	237	18.6 (6)				

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 34  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
5052	609	13.6 (107)	40.9 (16)	155	30.3 (78)	40.0 (28)	155	34.2 (19)	127	18.1 (5)
5054	80	*	*	38	*	*	19	*	31	*
5202	113	7.1 (8)	43.4 (33)	102	20.6 (18)	42.2 (42)	10	*	23	*
5203	57	*	*	52	*	*	9	*	17	*
5204	55	*	*	46	*	*	5	*	10	*
5205	76	*	*	75	*	*	8	*	24	*
5601	246	13.8 (108)	49.2 (90)	210	24.3 (46)	47.6 (79)	46	*	108	50.9 (43)
5801	759	16.2 (118)	43.6 (35)	644	22.2 (25)	46.3 (68)	126	20.6 (2)	116	25.0 (13)
5803	3	*	*	3	*	*	0	*	0	*
6103	1,440	10.6 (59)	48.5 (86)	1,332	23.0 (31)	48.3 (86)	238	35.3 (21)	163	43.6 (38)
6301	1,658	9.5 (31)	47.1 (69)	1,439	22.9 (30)	45.2 (63)	367	28.9 (12)	360	25.0 (13)
6312	993	12.1 (87)	44.3 (37)	879	24.6 (49)	46.8 (72)	244	30.3 (15)	283	29.7 (21)
6316	397	12.8 (96)	41.1 (20)	349	14.9 (2)	51.6 (109)	91	*	137	19.0 (7)
6320	87	*	*	78	*	*	19	*	28	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 35  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
6325	53	*	*	47	*	*	7	*	14	*	*	
6330	108	11.1 (68)	51.9 (110)	90	*	*	34	*	26	*	*	
6335	197	12.7 (93)	39.1 (11)	172	20.9 (20)	45.3 (64)	35	*	40	*	*	
6340	166	13.3(104)	41.0 (17)	145	17.2 (4)	47.6 (79)	29	*	35	*	*	
7304	97	*	*	79	*	*	12	*	26	*	*	
7306	971	7.3 (11)	53.9 (119)	853	21.2 (22)	51.0 (107)	163	35.6 (22)	255	47.8 (42)		
7308	70	*	*	58	*	*	11	*	16	*	*	
7401	784	15.7(117)	41.3 (23)	679	23.6 (38)	48.3 (86)	152	27.6 (8)	434	38.0 (34)		
7402	116	18.1(121)	36.2 (6)	102	33.3 (94)	45.1 (62)	36	*	63	*	*	
7403	36	*	*	33	*	*	12	*	20	*	*	
7404	47	*	*	43	*	*	10	*	19	*	*	
7601	226	8.4 (20)	46.9 (67)	204	23.0 (31)	44.1 (53)	44	*	74	*	*	
7850	200	11.5 (77)	48.0 (79)	176	29.0 (72)	42.0 (38)	57	*	101	30.7 (22)		
7851	38	*	*	31	*	*	10	*	14	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 36  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
7852	27	*	*	23	*	*	2	*	9	*	*	
8150	34	*	*	24	*	*	9	*	9	*	*	
8151	875	10.2 (42)	48.3 (83)	719	35.2 (99)	41.7 (36)	167	40.7 (25)	187	27.3 (16)		
8152	202	12.4 (91)	34.7 (4)	171	56.1 (115)	22.2 (4)	25	*	23	*	*	
8153	9	*	*	8	*	*	4	*	4	*	*	
8154	126	23.8 (125)	25.4 (2)	107	20.6 (18)	34.6 (14)	31	*	42	*	*	
8155	31	*	*	28	*	*	5	*	6	*	*	
8156	12	*	*	10	*	*	3	*	3	*	*	
8157	3	*	*	3	*	*	0	*	0	*	*	
8201	840	11.5 (77)	45.2 (52)	532	24.1 (43)	45.9 (66)	268	22.8 (4)	237	24.1 (12)		
8202	1,116	10.3 (48)	37.7 (8)	891	34.0 (95)	37.4 (20)	215	25.1 (7)	197	28.9 (19)		
8253	2	*	*	0	*	*	0	*	0	*	*	
8254	4	*	*	0	*	*	0	*	0	*	*	
8255	1,089	11.8 (82)	44.3 (37)	791	31.2 (84)	41.5 (34)	275	21.8 (3)	289	19.7 (9)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 37  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
8257	231	10.8 (63)	38.5 (9)	148	39.2 (105)	33.8 (11)	45	* *	41	* *
8258	550	10.7 (60)	44.9 (48)	425	24.7 (50)	48.9 (93)	122	37.7 (24)	91	* *
8901	376	10.1 (40)	50.5 (102)	312	32.4 (89)	38.8 (24)	15	* *	6	* *
8902	152	11.2 (71)	55.3 (120)	132	34.8 (98)	36.4 (18)	9	* *	14	* *
8903	434	6.0 (4)	56.2 (122)	355	36.1 (101)	42.8 (46)	41	* *	14	* *
8904	272	12.9 (98)	45.6 (57)	234	51.7 (113)	24.4 (5)	37	* *	49	* *
8905	723	8.6 (21)	49.7 (94)	584	42.8 (109)	31.3 (8)	13	* *	4	* *
9004	256	9.0 (26)	52.3 (112)	241	31.5 (86)	46.9 (74)	57	* *	83	* *
9010	76	* * * *	*	66	* * * *	*	9	* *	11	* *
9023	34	* * * *	*	30	* * * *	*	3	* *	1	* *
9040	174	10.9 (65)	46.6 (64)	154	27.9 (68)	48.1 (84)	48	* *	42	* *
9045	409	10.5 (53)	43.8 (36)	359	35.9 (100)	34.5 (13)	68	* *	53	* *
9050	740	9.6 (33)	45.0 (49)	634	30.1 (75)	40.9 (31)	97	* *	90	* *
9051	1,002	7.8 (13)	43.0 (31)	861	31.1 (83)	39.3 (27)	93	* *	345	32.2 (24)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 38  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators (2, 3)									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
9053	1,003	10.4 (50)	49.3 (92)	871	56.6 (116)	26.8 (7)	189	40.7 (25)	269	32.7 (25)
9054	178	7.9 (14)	53.4 (118)	155	30.3 (78)	46.5 (69)	31	* *	33	* *
9058	182	21.4 (124)	23.1 (1)	150	54.0 (114)	21.3 (3)	25	* *	18	* *
9066	905	10.2 (42)	51.6 (107)	782	34.3 (96)	42.1 (39)	145	40.7 (25)	222	54.5 (45)
9068	598	10.2 (42)	52.0 (111)	535	45.8 (112)	34.4 (12)	77	* *	87	* *
9069	770	8.2 (16)	50.9 (103)	696	34.3 (96)	42.7 (45)	163	33.7 (18)	312	35.3 (27)
9070	779	13.0 (100)	47.4 (72)	670	42.2 (107)	36.4 (18)	114	44.7 (29)	126	29.4 (20)
9072	569	9.3 (28)	49.4 (93)	501	36.7 (102)	40.1 (29)	84	* *	121	53.7 (44)
9074	520	8.3 (17)	51.5 (106)	464	25.2 (51)	49.6 (97)	101	34.7 (20)	100	37.0 (31)
9079	463	7.3 (11)	53.1 (117)	407	27.5 (66)	48.2 (85)	90	* *	95	* *
9080	766	6.8 (5)	48.0 (79)	642	27.4 (65)	44.4 (58)	128	42.2 (28)	103	35.0 (26)
9151	1	*	*	0	*	*	0	*	0	*
Unknown Clinics	328	7.0 (7)	47.9 (78)	260	42.7 (108)	35.4 (16)	71	* *	75	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**

Page: 39  
Run Date: 10 / 11 / 07

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number
<b>Michigan</b>	<b>58,427</b>	<b>10.9</b>	<b>46.3</b>	<b>48,821</b>	<b>29.3</b>	<b>43.6</b>	<b>10,047</b>	<b>30.5</b>	<b>14,170</b>
<b>Nation (Prior Year)</b>	<b>791,556</b>	<b>11.3</b>	<b>44.2</b>	<b>667,778</b>	<b>25.8</b>	<b>43.4</b>	<b>114,910</b>	<b>31.2</b>	<b>279,621</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 40  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
0101	52	* * * *		58	* *	
0102	137	89.8 (5)	2.2 (6)	158	44.3 (28)	
0103	150	86.7 (14)	4.7 (29)	168	39.9 (48)	
0104	50	* * * *		57	* *	
0250	50	* * * *		63	* *	
0251	40	* * * *		41	* *	
0252	43	* * * *		54	* *	
0253	45	* * * *		52	* *	
0401	174	90.8 (3)	2.3 (8)	194	39.7 (50)	
0402	152	92.1 (1)	2.0 (4)	171	47.4 (14)	
0403	58	* * * *		59	* *	
0404	50	* * * *		55	* *	
0501	186	79.0 (48)	12.9 (96)	201	47.3 (15)	
0502	2	* * * *		2	* *	
0503	63	* * * *		67	* *	
0504	217	88.9 (8)	5.5 (37)	238	45.8 (21)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 41  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
0505	142	83.8 (22)	9.2 (69)	154	50.0 (8)	
0506	65	* * * *		73	* *	
0650	345	80.3 (40)	5.5 (37)	379	34.8 (76)	
0651	216	81.0 (34)	5.1 (33)	250	43.2 (30)	
0652	114	71.9 (82)	15.8(102)	134	42.5 (35)	
0653	142	76.1 (64)	11.3 (82)	147	48.3 (11)	
0654	115	81.7 (32)	8.7 (64)	128	50.8 (5)	
0655	112	79.5 (44)	10.7 (79)	130	35.4 (74)	
0657	40	* * * *		42	* *	
0701	154	79.9 (42)	12.3 (92)	172	39.5 (53)	
0702	36	* * * *		38	* *	
0703	86	* * * *		93	* *	
0704	29	* * * *		32	* *	
0705	52	* * * *		59	* *	
0850	326	81.3 (33)	9.2 (69)	397	37.0 (65)	
0851	301	73.4 (78)	17.3(105)	334	47.0 (17)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 42  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
0901	603	81.8 (30)	7.3 (54)	672	42.9 (31)	
0902	19	* * * *		21	* *	
1051	474	64.8(109)	28.3(116)	538	50.4 (6)	
1101	574	69.0 (95)	11.8 (87)	720	34.6 (78)	
1102	288	84.7 (18)	2.1 (5)	335	30.4 (87)	
1106	43	* * * *		49	* *	
1250	340	70.6 (93)	17.9(106)	386	41.5 (38)	
1251	256	77.7 (54)	10.9 (81)	283	40.6 (44)	
1252	247	77.3 (57)	11.7 (85)	292	38.7 (60)	
1253	245	84.1 (20)	6.9 (49)	277	32.5 (82)	
1350	801	72.0 (80)	15.5(101)	902	39.4 (55)	
1351	49	* * * *		55	* *	
1352	134	62.7(112)	23.1(112)	152	40.8 (43)	
1401	91	* * * *		108	50.0 (8)	
1402	50	* * * *		60	* *	
1403	7	* * * *		8	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 43  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
1404	2	* * * *	*	8	*	*
1501	329	81.8 (30)	5.8 (40)	422	42.9 (31)	
1502	38	* * * *	*	44	*	*
1503	322	77.3 (57)	9.0 (67)	403	34.5 (79)	
1504	822	86.0 (15)	2.9 (11)	950	36.0 (70)	
1505	205	86.8 (13)	4.9 (31)	244	38.9 (57)	
1506	158	81.0 (34)	1.3 (1)	196	36.7 (66)	
1507	202	90.6 (4)	4.5 (27)	229	44.1 (29)	
1508	38	* * * *	*	85	*	*
1509	179	76.0 (65)	12.3 (92)	235	46.4 (19)	
1510	50	* * * *	*	70	*	*
1511	171	87.1 (11)	2.3 (8)	195	46.2 (20)	
1512	117	83.8 (22)	4.3 (24)	126	29.4 (88)	
1701	224	78.1 (51)	13.8 (97)	248	58.5 (1)	
1712	42	* * * *	*	44	*	*
1950	230	80.4 (39)	12.6 (94)	249	37.8 (63)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 44  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
1952	56	* * * *		58	* *	
1953	127	78.0 (52)	11.8 (87)	136	31.6 (84)	
1954	124	77.4 (56)	11.3 (82)	138	45.7 (22)	
1955	89	* * * *		95	* *	
1957	126	76.2 (62)	11.9 (89)	135	34.8 (76)	
1959	59	* * * *		66	* *	
1963	65	* * * *		67	* *	
2150	222	86.9 (12)	8.1 (62)	254	41.3 (41)	
2151	97	* * * *		115	42.6 (34)	
2251	166	77.1 (59)	15.1(100)	188	47.9 (13)	
2252	53	* * * *		66	* *	
2551	1,220	71.7 (87)	7.4 (55)	1,339	25.1 (98)	
2552	297	71.4 (90)	7.7 (56)	334	24.9 (99)	
2554	39	* * * *		45	* *	
2555	1,385	79.4 (46)	7.9 (60)	1,518	30.5 (86)	
3001	346	59.5(115)	20.5(110)	403	49.9 (10)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 45  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
3002	81	* * * *		87	* *	
3003	5	* * * *		7	* *	
3004	78	* * * *		89	* *	
3007	9	* * * *		12	* *	
3009	2	* * * *		3	* *	
3201	223	89.7 (6)	4.0 (20)	238	44.5 (27)	
3202	0	* * * *		1	* *	
3301	1,274	79.7 (43)	5.2 (34)	1,407	23.8(102)	
3302	130	77.7 (54)	3.1 (13)	143	21.0(111)	
3303	20	* * * *		22	* *	
3304	47	* * * *		55	* *	
3305	28	* * * *		31	* *	
3306	44	* * * *		52	* *	
3308	17	* * * *		17	* *	
3309	12	* * * *		15	* *	
3311	7	* * * *		6	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 46  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
3550	26	* * * *		30	* *	
3551	10	* * * *		9	* *	
3552	12	* * * *		14	* *	
3650	1	* * * *		7	* *	
3701	241	87.6 (9)	3.3 (15)	271	45.0 (24)	
3702	77	* * * *		87	* *	
3703	34	* * * *		34	* *	
3704	34	* * * *		40	* *	
3805	990	82.6 (26)	7.0 (50)	1,114	39.9 (48)	
3950	579	75.5 (72)	5.9 (43)	706	41.4 (39)	
3952	17	* * * *		22	* *	
3953	28	* * * *		31	* *	
3954	65	* * * *		77	* *	
4001	285	76.8 (61)	7.7 (56)	354	44.6 (26)	
4002	270	74.1 (77)	5.2 (34)	320	42.2 (37)	
4103	10	* * * *		12	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 47  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
4106	522	62.3(114)	24.1(114)	574	35.7 (72)	
4107	992	64.9(108)	25.1(115)	1,099	41.2 (42)	
4109	368	68.8 (96)	23.1(112)	396	39.6 (51)	
4110	577	53.6(117)	37.1(119)	648	51.4 (4)	
4111	15	* * * *		17	* *	
4112	243	50.6(119)	10.7 (79)	281	25.3 (96)	
4113	293	28.3(121)	49.1(120)	320	35.3 (75)	
4120	675	71.6 (89)	19.4(108)	729	36.6 (67)	
4160	705	71.9 (82)	16.0(103)	780	37.4 (64)	
4351	237	84.0 (21)	4.6 (28)	258	40.3 (46)	
4352	67	* * * *		75	* *	
4353	157	80.9 (37)	10.2 (74)	186	53.8 (3)	
4354	267	89.5 (7)	3.0 (12)	306	42.5 (35)	
4355	123	69.1 (94)	20.3(109)	131	38.9 (57)	
4356	205	79.5 (44)	11.7 (85)	217	39.2 (56)	
4357	0	* * * *		2	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 48  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
4358	65	* * * *		88	* *	
4359	127	81.9 (29)	7.9 (60)	135	48.1 (12)	
4360	71	* * * *		77	* *	
4361	89	* * * *		104	36.5 (68)	
4362	126	77.8 (53)	12.7 (95)	129	45.0 (24)	
4363	306	75.8 (69)	17.0(104)	333	41.4 (39)	
4401	230	54.8(116)	37.0(118)	445	40.0 (47)	
4501	314	85.0 (16)	4.1 (22)	344	38.4 (62)	
4601	506	84.8 (17)	2.4 (10)	552	26.3 (93)	
4701	354	75.7 (70)	11.3 (82)	385	22.9(106)	
5050	320	84.4 (19)	3.8 (19)	410	32.2 (83)	
5051	859	83.6 (24)	3.4 (17)	1,055	27.5 (92)	
5052	517	82.2 (27)	3.5 (18)	631	27.7 (90)	
5054	66	* * * *		87	* *	
5202	110	80.9 (37)	9.1 (68)	125	58.4 (2)	
5203	56	* * * *		60	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 49  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
5204	47	* * * *		55	*	*
5205	78	* * * *		81	*	*
5601	340	87.6 (9)	4.4 (25)	377	38.5 (61)	
5801	691	75.1 (73)	14.8 (99)	775	38.8 (59)	
5803	3	* * * *		3	*	*
6103	1,413	79.1 (47)	10.5 (76)	1,576	39.5 (53)	
6301	1,529	72.9 (79)	8.8 (65)	1,711	26.1 (95)	
6312	928	76.0 (65)	7.0 (50)	1,026	20.4(112)	
6316	363	77.1 (59)	10.5 (76)	401	24.7(100)	
6320	85	* * * *		91	*	*
6325	50	* * * *		53	*	*
6330	94	* * * *		108	20.4(112)	
6335	181	76.2 (62)	10.5 (76)	209	33.5 (81)	
6340	147	74.8 (75)	10.2 (74)	166	40.4 (45)	
7304	87	* * * *		99	*	*
7306	906	72.0 (80)	18.2(107)	983	39.6 (51)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 50  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
7308	61	* * * *		69	* *	
7401	740	68.2 (98)	22.8(111)	795	36.0 (70)	
7402	106	82.1 (28)	6.6 (47)	116	27.6 (91)	
7403	36	* * * *		37	* *	
7404	46	* * * *		48	* *	
7601	205	81.0 (34)	9.8 (73)	233	45.1 (23)	
7850	386	36.3(120)	58.8(121)	435	50.1 (7)	
7851	52	* * * *		58	* *	
7852	31	* * * *		33	* *	
8150	28	* * * *		36	* *	
8151	792	82.8 (25)	3.2 (14)	905	28.2 (89)	
8152	181	71.8 (85)	7.7 (56)	212	42.9 (31)	
8153	8	* * * *		9	* *	
8154	110	91.8 (2)	1.8 (2)	134	24.6(101)	
8155	27	* * * *		32	* *	
8156	11	* * * *		12	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 51  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
8157	3	* * * *	*	3	*	*
8201	766	78.6 (50)	4.4 (25)	861	18.4(119)	
8202	1,004	71.2 (91)	14.4 (98)	1,160	36.2 (69)	
8253	0	* * * *	*	2	*	*
8254	0	* * * *	*	4	*	*
8255	1,007	78.9 (49)	5.8 (40)	1,127	25.2 (97)	
8257	210	71.9 (82)	9.5 (72)	239	23.4(104)	
8258	514	75.1 (73)	7.8 (59)	561	21.2(110)	
8901	340	62.6(113)	7.1 (53)	383	19.6(116)	
8902	140	67.9 (99)	6.4 (45)	168	22.0(108)	
8903	377	65.3(106)	6.6 (47)	461	20.0(114)	
8904	246	75.6 (71)	4.9 (31)	282	26.2 (94)	
8905	614	75.9 (67)	6.5 (46)	781	34.3 (80)	
9004	250	66.0(105)	9.2 (69)	258	10.5(125)	
9010	72	* * * *	*	78	*	*
9023	36	* * * *	*	37	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 52  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
9040	170	75.9 (67)	1.8 (2)	175	16.6(121)	
9045	350	67.4(101)	12.0 (90)	420	23.8(102)	
9050	664	65.2(107)	12.0 (90)	769	35.6 (73)	
9051	869	53.3(118)	31.8(117)	1,037	47.2 (16)	
9053	916	66.9(102)	5.3 (36)	1,015	19.8(115)	
9054	165	66.7(104)	4.8 (30)	182	23.1(105)	
9058	170	80.0 (41)	8.8 (65)	186	46.8 (18)	
9066	808	71.7 (87)	2.2 (6)	913	18.6(118)	
9068	543	64.5(110)	5.7 (39)	612	21.6(109)	
9069	722	68.8 (96)	5.8 (40)	807	13.1(123)	
9070	730	66.8(103)	8.1 (62)	792	22.5(107)	
9072	523	67.5(100)	3.3 (15)	577	11.4(124)	
9074	497	63.8(111)	7.0 (50)	527	19.2(117)	
9079	448	74.3 (76)	4.2 (23)	470	14.0(122)	
9080	695	71.8 (85)	4.0 (20)	769	31.1 (85)	
9151	0	* * * *	*	1	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Page: 53  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)				
	Medical Care			WIC Enrollment	
	1st Trimester	No Care	1st Trimester	Number	% (Rank)
Unknown Clinics	290	71.0 (92)	5.9 (43)	373	17.4(120)
Michigan	<b>53,684</b>	<b>74.3</b>	<b>10.8</b>	<b>60,963</b>	<b>33.5</b>
Nation (Prior Year)	<b>498,811</b>	<b>77.6</b>	<b>8.2</b>	<b>722,757</b>	<b>32.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 54  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)					
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum	
	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0101	53 * *	44 * *	18 * *	0 * *	0 * *	0 * *
0102	142 60.6 (119)	121 32.2 (98)	58 * *	0 * *	0 * *	0 * *
0103	148 56.1 (112)	133 34.6 (107)	58 * *	0 * *	0 * *	0 * *
0104	55 * *	46 * *	23 * *	0 * *	0 * *	0 * *
0250	62 * *	50 * *	28 * *	0 * *	0 * *	0 * *
0251	41 * *	37 * *	17 * *	0 * *	0 * *	0 * *
0252	53 * *	40 * *	20 * *	0 * *	0 * *	0 * *
0253	49 * *	44 * *	24 * *	0 * *	0 * *	0 * *
0401	190 57.4 (115)	166 31.9 (96)	86 * *	0 * *	0 * *	0 * *
0402	169 53.8 (104)	143 30.8 (89)	62 * *	0 * *	0 * *	0 * *
0403	59 * *	54 * *	24 * *	0 * *	0 * *	0 * *
0404	52 * *	50 * *	21 * *	0 * *	0 * *	0 * *
0501	196 54.6 (107)	180 28.3 (78)	91 * *	0 * *	0 * *	0 * *
0502	2 * *	2 * *	0 * *	0 * *	0 * *	0 * *
0503	62 * *	60 * *	26 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 55  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)					
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum	
	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0504	227 47.6 (75)	183 29.0 (81)	78 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0505	147 51.7 (96)	126 29.4 (83)	62 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0506	70 * * * * * *	61 * * * * * *	25 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0650	313 55.9 (111)	242 36.8 (113)	102 22.5 (38)	0 * * * * * *	0 * * * * * *	0 * * * * * *
0651	198 73.7 (122)	178 42.1 (115)	92 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0652	102 56.9 (113)	87 * * * * * *	35 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0653	135 51.9 (97)	131 26.7 (69)	55 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0654	80 * * * * * *	54 * * * * * *	29 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0655	126 57.1 (114)	110 32.7 (100)	51 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0657	42 * * * * * *	36 * * * * * *	21 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0701	163 49.1 (82)	133 27.1 (71)	56 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0702	37 * * * * * *	34 * * * * * *	20 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0703	86 * * * * * *	82 * * * * * *	43 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0704	30 * * * * * *	28 * * * * * *	13 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0705	57 * * * * * *	51 * * * * * *	20 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 56  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)					
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum	
	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0850	313 55.0 (108)	255 32.2 (98)	94 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0851	289 50.9 (92)	210 29.5 (84)	71 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
0901	638 51.6 (95)	572 31.3 (92)	254 29.1 (31)	0 * * * * * *	0 * * * * * *	0 * * * * * *
0902	21 * * * * * *	16 * * * * * *	3 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
1051	524 46.4 (70)	446 23.3 (54)	177 33.9 (13)	0 * * * * * *	0 * * * * * *	0 * * * * * *
1101	691 33.3 (40)	590 18.6 (41)	164 29.3 (29)	0 * * * * * *	0 * * * * * *	0 * * * * * *
1102	317 49.8 (88)	266 33.8 (104)	122 25.4 (36)	0 * * * * * *	0 * * * * * *	0 * * * * * *
1106	45 * * * * * *	41 * * * * * *	23 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
1250	355 43.7 (58)	293 27.3 (72)	109 24.8 (37)	0 * * * * * *	0 * * * * * *	0 * * * * * *
1251	209 67.5 (121)	212 41.5 (114)	106 19.8 (39)	0 * * * * * *	0 * * * * * *	0 * * * * * *
1252	272 33.8 (42)	213 21.1 (46)	49 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
1253	267 43.4 (57)	213 29.1 (82)	73 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
1350	873 46.0 (67)	766 27.9 (75)	302 26.5 (35)	0 * * * * * *	0 * * * * * *	0 * * * * * *
1351	54 * * * * * *	47 * * * * * *	18 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *
1352	147 46.9 (72)	132 26.5 (68)	55 * * * * * *	0 * * * * * *	0 * * * * * *	0 * * * * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 57  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)					
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum	
	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
1401	107 49.5 (85)	85 * *	36 * *	0 * *	0 * *	0 * *
1402	59 * *	49 * *	15 * *	0 * *	0 * *	0 * *
1403	8 * *	6 * *	3 * *	0 * *	0 * *	0 * *
1404	6 * *	2 * *	0 * *	0 * *	0 * *	0 * *
1501	391 36.6 (49)	338 18.0 (38)	106 29.2 (30)	0 * *	0 * *	0 * *
1502	44 * *	34 * *	13 * *	0 * *	0 * *	0 * *
1503	355 34.9 (44)	262 23.3 (54)	78 * *	0 * *	0 * *	0 * *
1504	867 28.4 (29)	752 11.7 (23)	160 48.1 (1)	0 * *	0 * *	0 * *
1505	237 37.6 (50)	213 15.5 (30)	67 * *	0 * *	0 * *	0 * *
1506	179 52.5 (99)	169 33.7 (103)	68 * *	0 * *	0 * *	0 * *
1507	220 49.1 (82)	199 25.1 (63)	91 * *	0 * *	0 * *	0 * *
1508	52 * *	49 * *	0 * *	0 * *	0 * *	0 * *
1509	207 30.9 (33)	177 14.7 (28)	44 * *	0 * *	0 * *	0 * *
1510	65 * *	55 * *	22 * *	0 * *	0 * *	0 * *
1511	184 53.8 (104)	165 36.4 (112)	72 * *	0 * *	0 * *	0 * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 58  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1512	104	51.0 (94)	99	* *	31	* *	0	* *	0	* *
1701	247	50.2 (89)	212	31.6 (94)	99	* *	0	* *	0	* *
1712	44	* *	36	* *	18	* *	0	* *	0	* *
1950	244	49.6 (86)	205	34.1 (105)	90	* *	0	* *	0	* *
1952	54	* *	52	* *	19	* *	0	* *	0	* *
1953	135	49.6 (86)	121	29.8 (86)	44	* *	0	* *	0	* *
1954	138	47.1 (73)	114	33.3 (101)	47	* *	0	* *	0	* *
1955	94	* *	81	* *	34	* *	0	* *	0	* *
1957	111	54.1 (106)	113	27.4 (73)	37	* *	0	* *	0	* *
1959	53	* *	55	* *	20	* *	0	* *	0	* *
1963	64	* *	51	* *	17	* *	0	* *	0	* *
2150	240	55.4 (109)	199	25.6 (65)	88	* *	0	* *	0	* *
2151	93	* *	95	* *	37	* *	0	* *	0	* *
2251	171	49.1 (82)	162	29.6 (85)	66	* *	0	* *	0	* *
2252	65	* *	54	* *	16	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 59  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal			Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2551	1,242	35.8 (46)	1,080	21.0 (45)	303	31.4 (22)	0	*	*	0	*	*
2552	312	33.3 (40)	270	20.4 (43)	75	*	0	*	*	0	*	*
2554	44	*	38	*	11	*	0	*	*	0	*	*
2555	1,330	46.2 (68)	1,157	28.3 (78)	392	28.3 (34)	0	*	*	0	*	*
3001	385	32.5 (39)	316	22.5 (52)	94	*	0	*	*	0	*	*
3002	84	*	79	*	22	*	0	*	*	0	*	*
3003	7	*	5	*	0	*	0	*	*	0	*	*
3004	88	*	74	*	2	*	0	*	*	0	*	*
3007	11	*	8	*	1	*	0	*	*	0	*	*
3009	3	*	2	*	0	*	0	*	*	0	*	*
3201	231	42.0 (55)	204	21.6 (47)	78	*	0	*	*	0	*	*
3202	0	*	0	*	0	*	0	*	*	0	*	*
3301	1,248	42.5 (56)	1,074	23.0 (53)	337	31.5 (20)	0	*	*	0	*	*
3302	126	31.7 (35)	116	22.4 (51)	26	*	0	*	*	0	*	*
3303	17	*	15	*	6	*	0	*	*	0	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 60  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
3304	51	*	*	42	*	*	4	*	*	0	*	*
3305	28	*	*	30	*	*	8	*	*	0	*	*
3306	49	*	*	39	*	*	8	*	*	0	*	*
3308	15	*	*	14	*	*	7	*	*	0	*	*
3309	10	*	*	9	*	*	3	*	*	0	*	*
3311	6	*	*	7	*	*	4	*	*	0	*	*
3550	29	*	*	24	*	*	15	*	*	0	*	*
3551	9	*	*	9	*	*	3	*	*	0	*	*
3552	14	*	*	11	*	*	6	*	*	0	*	*
3650	7	*	*	1	*	*	1	*	*	0	*	*
3701	266	44.7	(63)	231	28.1	(76)	90	*	*	0	*	*
3702	87	*	*	70	*	*	35	*	*	0	*	*
3703	36	*	*	34	*	*	13	*	*	0	*	*
3704	38	*	*	30	*	*	11	*	*	0	*	*
3805	1,086	53.0	(101)	926	31.4	(93)	416	30.0	(28)	0	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 61  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)							
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3950	680	44.3 (62)	598	27.6 (74)	236	28.4 (33)	0	* * * * *
3952	21	* * *	19	* *	6	* * *	0	* * * * *
3953	31	* * *	28	* *	7	* * *	0	* * * * *
3954	73	* * *	65	* *	21	* * *	0	* * * * *
4001	343	35.9 (47)	265	23.4 (57)	74	* * *	0	* * * * *
4002	293	44.0 (61)	231	28.6 (80)	84	* * *	0	* * * * *
4103	12	* * *	10	* *	6	* * *	0	* * * * *
4106	567	25.4 (24)	497	14.9 (29)	104	32.7 (15)	0	* * * * *
4107	1,028	41.3 (52)	887	21.9 (48)	300	34.7 (12)	0	* * * * *
4109	386	44.8 (64)	352	25.0 (60)	131	30.5 (26)	0	* * * * *
4110	632	29.1 (30)	542	16.1 (33)	131	32.1 (18)	0	* * * * *
4111	17	* * *	14	* *	6	* * *	0	* * * * *
4112	280	1.8 (1)	256	0.0 (1)	5	* * *	0	* * * * *
4113	314	10.5 (4)	280	4.6 (5)	24	* * *	0	* * * * *
4120	712	36.2 (48)	638	19.3 (42)	188	36.7 (10)	0	* * * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 62  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4160	727	29.3 (32)	629	13.7 (27)	154	46.8 (2)	0	*	*	0	*	*
4351	251	49.0 (81)	224	34.8 (108)	94	*	0	*	*	0	*	*
4352	70	*	56	*	29	*	0	*	*	0	*	*
4353	181	48.1 (78)	148	25.0 (60)	68	*	0	*	*	0	*	*
4354	299	50.8 (91)	258	35.3 (109)	112	28.6 (32)	0	*	*	0	*	*
4355	126	61.1 (120)	116	31.9 (96)	58	*	0	*	*	0	*	*
4356	213	47.9 (76)	198	22.2 (49)	78	*	0	*	*	0	*	*
4357	1	*	0	*	0	*	0	*	*	0	*	*
4358	74	*	66	*	4	*	0	*	*	0	*	*
4359	133	35.3 (45)	120	23.3 (54)	31	*	0	*	*	0	*	*
4360	77	*	71	*	22	*	0	*	*	0	*	*
4361	100	60.0 (118)	85	*	46	*	0	*	*	0	*	*
4362	132	48.5 (80)	126	33.3 (101)	56	*	0	*	*	0	*	*
4363	333	50.5 (90)	295	29.8 (86)	130	32.3 (17)	0	*	*	0	*	*
4401	170	48.2 (79)	368	31.0 (91)	46	*	0	*	*	0	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 63  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4501	332	52.7 (100)	309	30.1 (88)	136	31.6 (19)	0	* * * * * *	0	* * * * * *
4601	548	47.1 (73)	478	24.9 (59)	184	41.8 (5)	0	* * * * * *	0	* * * * * *
4701	360	53.6 (103)	334	26.9 (70)	155	43.9 (3)	0	* * * * * *	0	* * * * * *
5050	380	47.9 (76)	65	* * * *	16	* * * *	0	* * * * * *	0	* * * * * *
5051	931	39.4 (51)	262	12.6 (25)	51	* * * *	0	* * * * * *	0	* * * * * *
5052	576	41.5 (53)	136	18.4 (40)	34	* * * *	0	* * * * * *	0	* * * * * *
5054	76	* * * *	32	* * * *	3	* * * *	0	* * * * * *	0	* * * * * *
5202	118	46.6 (71)	107	26.2 (67)	46	* * * *	0	* * * * * *	0	* * * * * *
5203	58	* * * *	53	* * * *	26	* * * *	0	* * * * * *	0	* * * * * *
5204	54	* * * *	44	* * * *	20	* * * *	0	* * * * * *	0	* * * * * *
5205	78	* * * *	74	* * * *	28	* * * *	0	* * * * * *	0	* * * * * *
5601	357	46.2 (68)	320	25.3 (64)	126	34.9 (11)	0	* * * * * *	0	* * * * * *
5801	765	53.3 (102)	645	30.9 (90)	287	31.4 (22)	0	* * * * * *	0	* * * * * *
5803	3	* * * *	3	* * * *	2	* * * *	0	* * * * * *	0	* * * * * *
6103	1,514	43.9 (60)	1,347	25.9 (66)	477	31.4 (22)	0	* * * * * *	0	* * * * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 64  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6301	1,641	31.1 (34)	1,438	16.6 (34)	361	40.4 (7)	0	* * *	0	* * *
6312	987	22.1 (22)	865	10.8 (20)	157	42.7 (4)	0	* * *	0	* * *
6316	391	43.7 (58)	351	28.2 (77)	119	30.3 (27)	0	* * *	0	* * *
6320	90	* * *	68	* * *	7	* * *	0	* * *	0	* * *
6325	53	* * *	45	* * *	23	* * *	0	* * *	0	* * *
6330	100	34.0 (43)	85	* * *	20	* * *	0	* * *	0	* * *
6335	197	32.0 (37)	172	16.9 (36)	49	* * *	0	* * *	0	* * *
6340	161	32.3 (38)	139	20.9 (44)	39	* * *	0	* * *	0	* * *
7304	97	* * *	75	* * *	20	* * *	0	* * *	0	* * *
7306	944	29.1 (30)	761	16.8 (35)	182	41.8 (5)	0	* * *	0	* * *
7308	67	* * *	51	* * *	20	* * *	0	* * *	0	* * *
7401	787	57.6 (117)	681	35.5 (110)	371	31.5 (20)	0	* * *	0	* * *
7402	115	57.4 (115)	101	35.6 (111)	50	* * *	0	* * *	0	* * *
7403	36	* * *	33	* * *	13	* * *	0	* * *	0	* * *
7404	48	* * *	44	* * *	23	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 65  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)					
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum	
	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
7601	228 55.7 (110)	209 31.6 (94)	105 33.3 (14)	0 * * *	0 * * *	0 * * *
7850	195 52.3 (98)	376 34.3 (106)	60 * * *	0 * * *	0 * * *	0 * * *
7851	37 * * *	51 * * *	18 * * *	0 * * *	0 * * *	0 * * *
7852	27 * * *	30 * * *	14 * * *	0 * * *	0 * * *	0 * * *
8150	35 * * *	27 * * *	8 * * *	0 * * *	0 * * *	0 * * *
8151	872 31.8 (36)	723 17.8 (37)	158 38.6 (8)	0 * * *	0 * * *	0 * * *
8152	201 44.8 (64)	180 22.2 (49)	65 * * *	0 * * *	0 * * *	0 * * *
8153	9 * * *	7 * * *	2 * * *	0 * * *	0 * * *	0 * * *
8154	119 14.3 (8)	97 * * *	13 * * *	0 * * *	0 * * *	0 * * *
8155	28 * * *	28 * * *	1 * * *	0 * * *	0 * * *	0 * * *
8156	10 * * *	9 * * *	4 * * *	0 * * *	0 * * *	0 * * *
8157	3 * * *	3 * * *	0 * * *	0 * * *	0 * * *	0 * * *
8201	809 41.5 (53)	511 18.2 (39)	151 37.7 (9)	0 * * *	0 * * *	0 * * *
8202	775 13.0 (7)	746 4.3 (4)	53 * * *	0 * * *	0 * * *	0 * * *
8253	2 * * *	0 * * *	0 * * *	0 * * *	0 * * *	0 * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 66  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)					
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household		
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum	
	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
8254	4 *	*	0 *	*	0 *	*
8255	858 50.9 (92)	658 24.3 (58)	230 31.3 (25)	0 *	*	*
8257	89 *	*	78 *	*	0 *	*
8258	473 45.5 (66)	352 25.0 (60)	95 *	*	0 *	*
8901	354 21.2 (20)	287 7.3 (11)	39 *	*	0 *	*
8902	152 17.1 (11)	126 9.5 (14)	18 *	*	0 *	*
8903	398 17.1 (11)	346 10.4 (17)	37 *	*	0 *	*
8904	262 16.4 (10)	225 6.7 (9)	30 *	*	0 *	*
8905	651 11.2 (5)	568 6.3 (6)	41 *	*	0 *	*
9004	245 18.4 (16)	234 9.4 (13)	24 *	*	0 *	*
9010	74 *	*	64 *	*	0 *	*
9023	30 *	*	29 *	*	3 *	*
9040	170 25.3 (23)	150 13.3 (26)	34 *	*	0 *	*
9045	364 10.2 (3)	310 6.5 (8)	24 *	*	0 *	*
9050	731 15.0 (9)	655 7.0 (10)	80 *	*	0 *	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 67  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household					
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum			
Clinic	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
9051	646	17.2	(13)	537	6.3	(6)	51	*	*	0	*	*
9053	827	26.6	(28)	654	15.9	(32)	123	32.5	(16)	0	*	*
9054	176	18.2	(15)	160	12.5	(24)	23	*	*	0	*	*
9058	160	7.5	(2)	152	2.6	(2)	6	*	*	0	*	*
9066	599	20.9	(19)	664	10.5	(18)	51	*	*	0	*	*
9068	430	26.0	(25)	406	10.3	(16)	58	*	*	0	*	*
9069	724	22.0	(21)	598	10.7	(19)	97	*	*	0	*	*
9070	607	20.6	(17)	541	9.6	(15)	67	*	*	0	*	*
9072	549	20.8	(18)	497	8.0	(12)	70	*	*	0	*	*
9074	507	26.0	(25)	462	11.0	(22)	91	*	*	0	*	*
9079	451	11.5	(6)	410	4.1	(3)	36	*	*	0	*	*
9080	763	18.1	(14)	614	10.9	(21)	73	*	*	0	*	*
9151	1	*	*	0	*	*	0	*	*	0	*	*
Unknown Clinics	301	26.2	(27)	232	15.5	(30)	47	*	*	0	*	*
<b>Michigan</b>	<b>54,944</b>	<b>37.6</b>		<b>46,663</b>	<b>21.1</b>		<b>13,705</b>	<b>32.8</b>		<b>0</b>	<b>*</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**

Page: 68  
Run Date: 10 / 11 / 07

<b>Comparison of Smoking Indicators</b> (2, 3)							
<b>Clinic</b>	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household			
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum		
	Number (Rank)	Number (Rank)	Number (Rank)	Number (Rank)	Number (Rank)		
<b>Nation (Prior Year)</b>	<b>677,683</b> 30.0	<b>553,373</b> 19.2	<b>140,038</b> 33.0	<b>362,509</b> 28.5	<b>443,894</b> 26.1		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 69  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0101	41	* * * *	45	* *	41	* *	44	*	*	*
0102	120	4.2 (11) 16.7 (118)	131	0.0 (1)	120	4.2 (11)	124	46.8 (90)		
0103	135	5.2 (27) 8.9 (72)	145	0.7 (115)	134	4.5 (14)	143	44.1 (95)		
0104	45	* * * *	50	* *	44	* *	49	*	*	*
0250	47	* * * *	50	* *	47	* *	49	*	*	*
0251	36	* * * *	37	* *	36	* *	37	*	*	*
0252	42	* * * *	42	* *	41	* *	39	*	*	*
0253	42	* * * *	43	* *	42	* *	39	*	*	*
0401	162	9.9 (98) 13.0 (114)	164	0.0 (1)	162	9.9 (99)	158	51.3 (74)		
0402	144	5.6 (31) 8.3 (66)	144	0.0 (1)	143	5.6 (31)	141	70.9 (11)		
0403	56	* * * *	56	* *	55	* *	56	*	*	*
0404	47	* * * *	50	* *	47	* *	46	*	*	*
0501	177	4.5 (15) 12.4 (111)	183	0.0 (1)	176	4.5 (14)	161	71.4 (10)		
0502	2	* * * *	2	* *	2	* *	2	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 70  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0503	57	* * * *	64	* *	57	* *	63	* *		
0504	204	5.9 (35) 10.3 (97)	215	0.0 (1)	204	5.9 (37)	190	79.5 (6)		
0505	132	6.8 (53) 9.1 (78)	140	0.0 (1)	132	6.8 (51)	131	69.5 (13)		
0506	60	* * * *	62	* *	60	* *	61	* *		
0650	314	7.0 (59) 7.3 (52)	339	0.0 (1)	313	7.0 (59)	318	70.8 (12)		
0651	201	8.5 (84) 6.0 (34)	212	0.0 (1)	201	8.5 (83)	186	51.1 (75)		
0652	108	4.6 (17) 11.1 (104)	112	0.0 (1)	106	4.7 (20)	108	52.8 (67)		
0653	120	5.0 (22) 10.0 (94)	133	0.0 (1)	120	5.0 (23)	131	61.1 (36)		
0654	91	* * * *	95	* *	91	* *	87	* *		
0655	109	7.3 (64) 9.2 (80)	110	0.0 (1)	109	7.3 (65)	104	52.9 (65)		
0657	37	* * * *	37	* *	37	* *	37	* *		
0701	115	3.5 (6) 7.8 (59)	139	0.0 (1)	115	3.5 (6)	133	75.2 (7)		
0702	31	* * * *	33	* *	31	* *	33	* *		
0703	79	* * * *	79	* *	79	* *	70	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 71  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0704	27	* * * *	28	* *	27	* *	27	*	*	*
0705	43	* * * *	51	* *	43	* *	50	*	*	*
0850	312	7.7 (71) 10.9 (102)	340	0.3 (98)	311	7.7 (71)	320	64.1 (21)		
0851	285	6.0 (37) 9.5 (85)	297	0.3 (98)	284	6.0 (38)	288	50.0 (79)		
0901	591	6.3 (45) 8.1 (64)	600	0.0 (1)	589	6.3 (44)	581	49.4 (83)		
0902	17	* * * *	17	* *	17	* *	17	*	*	*
1051	448	6.9 (56) 10.7 (100)	460	0.0 (1)	447	6.9 (56)	442	79.6 (5)		
1101	594	8.6 (86) 7.4 (54)	597	0.0 (1)	591	8.6 (86)	571	49.2 (84)		
1102	273	4.4 (13) 9.2 (80)	281	0.0 (1)	273	4.4 (13)	265	56.2 (55)		
1106	43	* * * *	45	* *	43	* *	42	*	*	*
1250	326	6.1 (40) 6.1 (36)	338	0.0 (1)	326	6.1 (40)	330	58.5 (47)		
1251	248	3.2 (4) 6.9 (47)	252	0.0 (1)	248	3.2 (4)	244	57.4 (53)		
1252	203	4.4 (13) 5.4 (27)	245	0.4 (107)	202	4.5 (14)	233	58.8 (45)		
1253	215	8.4 (80) 9.3 (83)	233	0.0 (1)	214	8.4 (80)	226	42.5 (98)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 72  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
1350	742	6.6 (47) 10.2 (95)	783	0.0 (1)	740	6.6 (47)	712	50.4 (77)		
1351	46	* * * *	48	* *	45	* *	47	* *		
1352	128	4.7 (19) 3.9 (9)	132	0.0 (1)	127	4.7 (20)	127	31.5 (113)		
1401	86	* * * *	89	* *	85	* *	85	* *		
1402	52	* * * *	52	* *	52	* *	49	* *		
1403	7	* * * *	7	* *	7	* *	6	* *		
1404	2	* * * *	2	* *	2	* *	2	* *		
1501	321	5.0 (22) 8.4 (67)	345	0.0 (1)	320	5.0 (23)	331	63.4 (24)		
1502	36	* * * *	37	* *	36	* *	34	* *		
1503	313	7.0 (59) 8.0 (62)	330	0.0 (1)	313	7.0 (59)	311	61.1 (36)		
1504	702	9.8 (97) 8.8 (70)	806	0.2 (91)	699	9.7 (96)	777	62.5 (29)		
1505	211	5.7 (33) 12.8 (112)	215	0.0 (1)	211	5.7 (34)	210	61.0 (39)		
1506	167	3.0 (3) 12.0 (109)	170	0.6 (113)	167	3.0 (3)	166	63.9 (23)		
1507	166	4.2 (11) 14.5 (117)	200	0.0 (1)	165	4.2 (11)	198	61.1 (36)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 73  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
1508	19	* * * *	54	* *	19	* *	53	* *		
1509	142	2.8 (2) 9.2 (80)	184	0.5 (111)	141	2.8 (2)	178	60.7 (40)		
1510	52	* * * *	57	* *	51	* *	55	* *		
1511	164	5.5 (30) 12.2 (110)	167	0.0 (1)	164	5.5 (30)	162	59.3 (43)		
1512	107	5.6 (31) 6.5 (41)	108	0.0 (1)	107	5.6 (31)	104	52.9 (65)		
1701	211	5.7 (33) 12.8 (112)	213	0.0 (1)	211	5.7 (34)	207	63.3 (25)		
1712	36	* * * *	40	* *	36	* *	40	* *		
1950	214	7.0 (59) 4.7 (17)	216	0.0 (1)	211	7.1 (63)	197	58.4 (48)		
1952	52	* * * *	53	* *	50	* *	52	* *		
1953	120	6.7 (50) 10.8 (101)	124	1.6 (119)	116	6.9 (56)	120	55.8 (56)		
1954	118	5.1 (26) 10.2 (95)	113	0.9 (118)	112	4.5 (14)	105	62.9 (27)		
1955	82	* * * *	80	* *	79	* *	77	* *		
1957	117	8.5 (84) 11.1 (104)	119	0.0 (1)	117	8.5 (83)	115	66.1 (17)		
1959	52	* * * *	54	* *	52	* *	50	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 74  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)	Number (Rank)			
1963	54	* * * *	54	*	*	52	* *	53	*	*
2150	208	6.3 (45) 9.1 (78)	213	0.0 (1)		208	6.3 (44)	195	62.1 (32)	
2151	97	* * * *	98	*	*	97	* *	87	*	*
2251	165	6.7 (50) 6.1 (36)	165	0.0 (1)		165	6.7 (50)	164	59.8 (42)	
2252	53	* * * *	52	*	*	52	* *	51	*	*
2551	1,130	9.3 (92) 4.0 (11)	1,147	0.0 (1)		1,127	9.3 (92)	1,097	36.6 (105)	
2552	263	8.0 (75) 4.6 (15)	281	0.4 (107)		262	8.0 (75)	279	40.5 (100)	
2554	34	* * * *	39	*	*	34	* *	36	*	*
2555	1,181	7.5 (70) 8.5 (68)	1,287	0.2 (91)		1,177	7.6 (70)	1,269	49.5 (82)	
3001	331	13.0 (112) 4.8 (20)	331	0.0 (1)		330	13.0 (112)	309	34.6 (107)	
3002	81	* * * *	80	*	*	80	* *	74	*	*
3003	5	* * * *	5	*	*	5	* *	5	*	*
3004	74	* * * *	74	*	*	74	* *	71	*	*
3007	8	* * * *	8	*	*	8	* *	8	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 75  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
3009	2	* * * *	2	*	2	*	2	*	*	*
3201	204	7.4 (68) 11.8 (108)	206	0.0 (1)	203	7.4 (68)	200	45.0 (91)		
3202	0	* * * *	0	*	0	*	0	*	*	*
3301	1,071	7.0 (59) 7.8 (59)	1,211	0.0 (1)	1,069	6.9 (56)	1,119	64.1 (21)		
3302	115	10.4 (103) 3.5 (4)	124	0.0 (1)	115	10.4 (103)	116	58.6 (46)		
3303	19	* * * *	20	*	19	*	18	*	*	*
3304	42	* * * *	45	*	42	*	45	*	*	*
3305	27	* * * *	30	*	27	*	30	*	*	*
3306	41	* * * *	46	*	41	*	43	*	*	*
3308	13	* * * *	16	*	13	*	16	*	*	*
3309	12	* * * *	12	*	12	*	11	*	*	*
3311	6	* * * *	6	*	5	*	7	*	*	*
3550	25	* * * *	25	*	25	*	25	*	*	*
3551	10	* * * *	10	*	10	*	10	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 76  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>							
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed
	Low (4)	High (5)	Number (Rank)	Number (Rank)	Number (Rank)	Number (Rank)	Number (Rank)	
3552	11	* * * *	11	* *	11	* *	11	* *
3650	1	* * * *	1	* *	1	* *	1	* *
3701	232	3.4 (5) 11.2 (106)	233	0.0 (1)	232	3.4 (5)	223	48.9 (85)
3702	71	* * * *	72	* *	71	* *	71	* *
3703	34	* * * *	34	* *	34	* *	33	* *
3704	30	* * * *	30	* *	30	* *	28	* *
3805	955	8.6 (86) 7.5 (56)	974	0.1 (83)	954	8.6 (86)	897	49.7 (80)
3950	604	8.1 (76) 10.3 (97)	609	0.0 (1)	604	8.1 (76)	589	55.7 (57)
3952	18	* * * *	18	* *	18	* *	18	* *
3953	28	* * * *	28	* *	28	* *	27	* *
3954	64	* * * *	65	* *	64	* *	65	* *
4001	259	9.3 (92) 4.6 (15)	265	0.4 (107)	259	9.3 (92)	246	54.5 (60)
4002	245	6.1 (40) 9.8 (92)	256	0.0 (1)	244	6.1 (40)	242	61.6 (34)
4103	10	* * * *	10	* *	10	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 77  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>										
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed			
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	
4106	500	8.4 (80)	7.0 (48)	501	0.0 (1)	500	8.4 (80)	440	52.5 (69)		
4107	884	9.5 (95)	9.6 (89)	933	0.1 (83)	879	9.4 (94)	876	51.7 (71)		
4109	350	9.4 (94)	9.7 (91)	352	0.0 (1)	349	9.5 (95)	334	62.3 (31)		
4110	531	4.5 (15)	8.9 (72)	541	0.0 (1)	530	4.5 (14)	513	60.0 (41)		
4111	14	*	*	*	*	14	*	*	11	*	*
4112	249	6.0 (37)	11.2 (106)	247	0.0 (1)	247	5.7 (34)	246	92.7 (3)		
4113	276	2.5 (1)	6.5 (41)	278	0.0 (1)	276	2.5 (1)	257	73.2 (8)		
4120	639	5.2 (27)	8.9 (72)	639	0.0 (1)	638	5.2 (29)	591	58.0 (51)		
4160	664	5.4 (29)	7.2 (50)	671	0.3 (98)	661	5.1 (28)	658	55.6 (58)		
4351	223	6.7 (50)	5.4 (27)	230	0.4 (107)	222	6.8 (51)	209	58.4 (48)		
4352	62	*	*	*	*	63	*	*	59	*	*
4353	149	8.7 (89)	4.7 (17)	149	0.0 (1)	148	8.8 (89)	139	61.2 (35)		
4354	254	9.1 (91)	13.8 (115)	261	0.0 (1)	253	9.1 (90)	254	63.0 (26)		
4355	114	3.5 (6)	8.8 (70)	114	0.0 (1)	114	3.5 (6)	105	64.8 (20)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 78  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
4356	197	6.1 (40) 5.6 (29)	195	0.0 (1)	195	6.2 (42)	189	54.5 (60)		
4357	0	* * * *	0	* *	0	* *	0	*	*	*
4358	64	* * * *	68	* *	64	* *	67	*	*	*
4359	116	6.9 (56) 9.5 (85)	127	0.0 (1)	115	7.0 (59)	104	62.5 (29)		
4360	69	* * * *	67	* *	67	* *	66	*	*	*
4361	85	* * * *	85	* *	85	* *	81	*	*	*
4362	124	4.0 (10) 6.5 (41)	124	0.0 (1)	123	4.1 (10)	121	69.4 (14)		
4363	289	5.9 (35) 14.2 (116)	291	0.0 (1)	287	5.6 (31)	280	66.8 (16)		
4401	355	9.0 (90) 9.0 (76)	377	0.5 (111)	353	9.1 (90)	365	48.5 (88)		
4501	310	6.8 (53) 10.6 (99)	309	0.0 (1)	309	6.8 (51)	303	58.1 (50)		
4601	475	8.2 (79) 8.2 (65)	477	0.0 (1)	475	8.2 (78)	468	53.0 (64)		
4701	333	3.6 (8) 9.0 (76)	336	0.0 (1)	329	3.6 (8)	325	69.2 (15)		
5050	83	* * * *	87	* *	83	* *	87	*	*	*
5051	329	4.6 (17) 9.4 (84)	339	0.0 (1)	328	4.6 (19)	337	94.4 (2)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 79  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>							
	Birthweight			Preterm (6)	Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number (Rank)		Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
5052	161	5.0 (22) 5.0 (24)	162	0.6 (113)	159	5.0 (23)	161	95.7 (1)
5054	38	* * * *	42	* *	38	* *	42	* *
5202	100	5.0 (22) 11.0 (103)	108	0.0 (1)	100	5.0 (23)	102	61.8 (33)
5203	50	* * * *	52	* *	49	* *	50	* *
5204	45	* * * *	45	* *	44	* *	43	* *
5205	74	* * * *	75	* *	73	* *	71	* *
5601	323	8.4 (80) 9.6 (89)	327	0.0 (1)	323	8.4 (80)	318	62.6 (28)
5801	640	6.6 (47) 6.1 (36)	647	0.0 (1)	637	6.6 (47)	605	48.8 (86)
5803	3	* * * *	3	* *	3	* *	3	* *
6103	1,342	6.6 (47) 6.8 (44)	1,357	0.1 (83)	1,337	6.6 (47)	1,261	44.6 (92)
6301	1,426	6.8 (53) 7.1 (49)	1,479	0.0 (1)	1,425	6.8 (51)	1,421	47.7 (89)
6312	874	8.6 (86) 9.8 (92)	901	0.0 (1)	873	8.6 (86)	870	52.1 (70)
6316	357	8.4 (80) 7.6 (57)	355	0.0 (1)	355	8.5 (83)	355	54.4 (62)
6320	78	* * * *	83	* *	75	* *	82	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 80  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
6325	46	* * * *	47	* *	45	* *	48	* *		
6330	85	* * * *	90	* *	84	* *	88	* *		
6335	161	6.2 (43) 6.2 (39)	184	0.0 (1)	161	6.2 (42)	183	49.7 (80)		
6340	137	3.6 (8) 8.0 (62)	145	0.0 (1)	135	3.7 (9)	140	55.0 (59)		
7304	80	* * * *	80	* *	80	* *	71	* *		
7306	859	9.9 (98) 6.8 (44)	860	0.1 (83)	856	9.8 (97)	764	44.1 (95)		
7308	57	* * * *	57	* *	57	* *	50	* *		
7401	679	8.1 (76) 4.9 (21)	681	0.0 (1)	679	8.1 (76)	673	43.5 (97)		
7402	99	* * * *	100	0.0 (1)	99	* *	98	* *		
7403	33	* * * *	33	* *	33	* *	33	* *		
7404	44	* * * *	44	* *	44	* *	44	* *		
7601	208	4.8 (20) 7.7 (58)	209	0.0 (1)	207	4.8 (22)	195	48.7 (87)		
7850	368	7.3 (64) 6.0 (34)	372	0.0 (1)	368	7.3 (65)	346	57.5 (52)		
7851	51	* * * *	51	* *	51	* *	45	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 81  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
7852	28	* * * *	29	* *	28	* *	29	* *		
8150	24	* * * *	26	* *	23	* *	27	* *		
8151	682	7.3 (64) 8.9 (72)	747	0.1 (83)	675	6.8 (51)	657	66.1 (17)		
8152	180	10.0 (101) 2.8 (3)	182	0.0 (1)	180	10.0 (100)	171	50.3 (78)		
8153	8	* * * *	8	* *	8	* *	8	* *		
8154	110	7.3 (64) 7.3 (52)	112	0.0 (1)	110	7.3 (65)	112	86.6 (4)		
8155	27	* * * *	29	* *	27	* *	24	* *		
8156	10	* * * *	10	* *	10	* *	9	* *		
8157	2	* * * *	3	* *	2	* *	2	* *		
8201	530	6.2 (43) 6.8 (44)	541	0.2 (91)	525	6.3 (44)	512	52.7 (68)		
8202	788	4.9 (21) 8.5 (68)	929	0.1 (83)	784	5.0 (23)	890	73.0 (9)		
8253	0	* * * *	0	* *	0	* *	0	* *		
8254	0	* * * *	0	* *	0	* *	0	* *		
8255	793	6.9 (56) 7.8 (59)	815	0.2 (91)	786	7.0 (59)	788	50.5 (76)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 82  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>							
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8257	135	7.4 (68) 4.4 (14)	156	0.0 (1)	135	7.4 (68)	153	65.4 (19)
8258	423	7.1 (63) 9.5 (85)	433	0.2 (91)	422	7.1 (63)	422	44.3 (93)
8901	306	11.1 (105) 5.6 (29)	322	0.3 (98)	304	11.2 (105)	297	34.3 (108)
8902	141	13.5 (114) 5.7 (31)	145	0.0 (1)	141	13.5 (114)	144	37.5 (104)
8903	366	10.1 (102) 4.1 (12)	378	0.3 (98)	366	10.1 (102)	358	28.2 (115)
8904	227	12.3 (109) 2.6 (2)	245	0.0 (1)	227	12.3 (109)	239	53.6 (63)
8905	598	7.9 (73) 9.5 (85)	629	0.3 (98)	594	7.9 (73)	599	34.1 (109)
9004	241	13.7 (115) 6.2 (39)	243	0.0 (1)	241	13.7 (115)	232	22.8 (118)
9010	61	* * * *	67	* *	61	* *	65	* *
9023	31	* * * *	31	* *	31	* *	31	* *
9040	153	7.8 (72) 3.9 (9)	156	0.0 (1)	153	7.8 (72)	154	26.0 (117)
9045	350	9.7 (96) 5.7 (31)	364	0.3 (98)	348	9.8 (97)	350	39.7 (101)
9050	618	7.9 (73) 4.7 (17)	657	0.0 (1)	617	7.9 (73)	642	51.7 (71)
9051	754	6.0 (37) 7.4 (54)	875	0.1 (83)	750	6.0 (38)	830	56.9 (54)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 83  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators <sup>(2, 3)</sup>							
	Birthweight		Preterm (6)	Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)		Number	% (Rank)	Number	% (Rank)	Number
9053	812	11.5(106)	5.0 (24)	877	0.0 (1)	812	11.5(106)	669
9054	161	9.9 (98)	3.7 (6)	160	0.0 (1)	160	10.0(100)	154
9058	148	8.1 (76)	2.0 (1)	155	0.0 (1)	147	8.2 (78)	154
9066	722	13.9(116)	3.5 (4)	792	0.1 (83)	721	13.9(116)	777
9068	520	12.3(109)	5.8 (33)	543	0.2 (91)	519	12.3(109)	425
9069	683	13.3(113)	5.1 (26)	722	0.0 (1)	683	13.3(113)	636
9070	630	12.2(108)	3.8 (7)	681	0.3 (98)	626	12.3(109)	647
9072	501	15.0(118)	3.8 (7)	510	0.8 (117)	496	15.1(118)	503
9074	466	14.2(117)	4.1 (12)	469	0.0 (1)	466	14.2(117)	421
9079	389	12.3(109)	4.9 (21)	413	0.2 (91)	386	12.2(108)	391
9080	643	10.7(104)	7.2 (50)	641	0.3 (98)	637	10.7(104)	640
9151	0	*	*	0	*	0	*	0
Unknown Clinics	246	11.8(107)	4.9 (21)	275	0.7 (115)	246	11.8(107)	262
								32.8(112)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**

Page: 84  
Run Date: 10 / 11 / 07

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Michigan	48,340	7.8	7.6	50,521	0.1	48,152	7.8	47,818	52.8
Nation (Prior Year)	707,018	8.3	7.0	636,104	12.4	583,078	3.8	704,334	59.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 85  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
0101	157	96.8	1.3	0.6	0.0	0.0	1.3
0102	499	95.2	1.8	1.4	0.0	0.4	1.2
0103	531	97.9	0.6	0.0	0.0	0.0	1.5
0104	169	97.6	0.0	1.8	0.0	0.0	0.6
0250	199	93.5	0.0	0.0	3.0	0.0	3.5
0251	150	96.7	0.0	0.0	2.0	0.0	1.3
0252	159	76.7	1.3	0.6	11.3	0.6	9.4
0253	182	87.9	0.0	0.0	10.4	0.0	1.6
0401	585	97.6	0.3	0.2	0.2	0.0	1.7
0402	553	92.8	0.5	1.1	0.4	0.5	4.7
0403	157	99.4	0.0	0.0	0.6	0.0	0.0
0404	178	94.9	0.0	2.2	0.6	0.6	1.7
0501	609	97.9	0.3	0.5	0.2	0.2	1.0
0502	3	*	*	*	*	*	*
0503	182	95.1	0.5	2.7	0.5	0.5	0.5
0504	701	89.9	0.7	1.1	4.1	0.6	3.6
0505	508	90.4	1.0	2.8	0.2	0.0	5.7
0506	229	97.4	0.4	1.3	0.0	0.0	0.9
0650	1,179	91.0	2.9	2.3	0.9	0.9	2.0
0651	745	97.6	0.5	0.7	0.1	0.3	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 86  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
0652	447	97.1	0.2	1.1	0.0	0.0	1.6
0653	479	98.5	0.0	0.2	0.0	0.0	1.3
0654	416	98.3	0.2	0.2	0.0	0.0	1.2
0655	481	97.9	0.2	0.0	0.0	0.4	1.5
0657	126	96.0	1.6	0.0	0.8	0.0	1.6
0701	511	90.8	1.8	0.8	0.4	5.5	0.8
0702	126	99.2	0.0	0.0	0.0	0.0	0.8
0703	276	93.8	1.4	1.1	0.7	1.4	1.4
0704	96	*	*	*	*	*	*
0705	189	97.4	0.0	0.0	0.0	0.5	2.1
0850	1,247	91.0	3.1	2.3	0.0	0.6	2.9
0851	960	93.8	0.5	3.8	0.1	0.6	1.3
0901	2,158	88.0	2.8	6.4	0.1	0.4	2.2
0902	84	*	*	*	*	*	*
1051	1,626	90.9	0.9	4.9	1.5	0.7	1.2
1059	4	*	*	*	*	*	*
1060	1	*	*	*	*	*	*
1101	2,148	37.3	56.9	3.4	0.2	1.5	0.7
1102	1,020	81.7	12.0	3.9	0.2	0.9	1.4
1106	120	90.8	0.8	2.5	0.0	0.0	5.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 87  
Run Date: 10 / 11 / 07

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan Native %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other/Unknown %</b>	
1250	1,139	88.4	0.5	9.4	0.0	0.5	1.1	
1251	916	98.1	0.5	0.8	0.0	0.2	0.3	
1252	945	64.9	1.1	33.2	0.1	0.1	0.6	
1253	879	87.5	9.1	2.5	0.1	0.0	0.8	
1350	2,870	66.4	21.8	8.9	0.4	1.1	1.4	
1351	212	95.8	0.0	2.8	0.0	0.0	1.4	
1352	551	62.6	31.0	4.4	0.0	0.2	1.8	
1401	325	88.3	1.5	4.6	0.6	0.3	4.6	
1402	193	56.0	0.0	19.7	21.2	0.5	2.6	
1403	30	*	*	*	*	*	*	
1404	22	*	*	*	*	*	*	
1405	1	*	*	*	*	*	*	
1501	1,369	55.7	8.7	33.8	0.2	0.2	1.3	
1502	118	80.5	0.8	15.3	0.0	0.0	3.4	
1503	1,359	53.8	5.4	37.1	0.5	0.4	2.7	
1504	2,882	45.5	3.5	44.0	0.1	5.7	1.2	
1505	700	86.0	1.1	9.0	0.3	2.0	1.6	
1506	564	85.5	0.9	11.2	0.0	0.0	2.5	
1507	707	94.1	1.1	2.5	0.1	0.4	1.7	
1508	303	0.7	0.0	99.0	0.0	0.0	0.3	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 88  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
1509	733	55.7	2.3	41.6	0.0	0.0	0.4
1510	264	90.5	4.2	4.5	0.0	0.0	0.8
1511	598	95.3	1.2	1.7	0.2	0.2	1.5
1512	332	93.7	0.9	2.7	0.3	0.3	2.1
1701	773	64.3	1.2	1.3	22.4	1.4	9.4
1712	138	65.2	0.0	0.0	23.2	1.4	10.1
1950	786	96.9	0.5	1.7	0.1	0.0	0.8
1952	186	98.4	0.0	0.0	0.0	0.0	1.6
1953	445	96.2	0.4	2.7	0.0	0.0	0.7
1954	446	92.2	0.9	6.1	0.0	0.0	0.9
1955	230	91.7	0.4	7.0	0.0	0.4	0.4
1956	14	*	*	*	*	*	*
1957	417	93.5	0.5	4.8	0.0	0.5	0.7
1959	229	93.9	3.1	2.2	0.0	0.0	0.9
1963	256	91.0	0.4	7.0	0.4	0.0	1.2
2150	830	94.6	0.4	0.1	0.5	0.7	3.7
2151	361	95.6	0.6	1.9	0.0	0.3	1.7
2251	606	96.7	0.7	0.5	0.3	1.3	0.5
2252	228	93.9	1.3	3.1	0.0	0.0	1.8
2551	4,419	32.1	64.7	1.4	0.1	0.2	1.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 89  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %	
2552	1,104	30.0	63.9	4.3	0.1	0.2	1.5	
2554	124	94.4	3.2	0.8	0.8	0.8	0.0	
2555	4,504	80.0	15.9	2.2	0.2	0.6	1.1	
3001	1,256	13.4	68.6	16.2	0.1	0.4	1.4	
3002	238	19.3	59.7	20.2	0.4	0.0	0.4	
3003	34	*	*	*	*	*	*	
3004	266	2.6	0.0	96.6	0.0	0.0	0.8	
3007	35	*	*	*	*	*	*	
3009	10	*	*	*	*	*	*	
3010	1	*	*	*	*	*	*	
3201	686	95.3	0.1	3.4	0.0	0.0	1.2	
3202	23	*	*	*	*	*	*	
3301	4,570	55.4	26.0	12.8	0.3	3.4	2.1	
3302	476	27.5	46.0	17.0	0.2	6.3	2.9	
3303	88	*	*	*	*	*	*	
3304	213	28.6	16.0	7.5	0.9	46.0	0.9	
3305	117	78.6	10.3	6.0	0.0	2.6	2.6	
3306	120	91.7	3.3	3.3	0.0	0.0	1.7	
3308	72	*	*	*	*	*	*	
3309	49	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 90  
Run Date: 10 / 11 / 07

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan Native %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other/Unknown %</b>	
3311	21	*	*	*	*	*	*	
3550	105	23.8	1.0	1.0	41.0	0.0	33.3	
3551	33	*	*	*	*	*	*	
3552	52	*	*	*	*	*	*	
3650	37	*	*	*	*	*	*	
3701	822	88.2	0.9	7.8	0.2	0.9	2.1	
3702	273	92.3	1.1	2.6	0.0	0.7	3.3	
3703	126	93.7	0.0	4.0	0.0	0.0	2.4	
3704	112	83.9	0.9	11.6	0.0	1.8	1.8	
3805	3,714	78.7	13.8	5.1	0.5	0.4	1.4	
3950	2,327	68.4	21.0	6.8	0.1	1.5	2.3	
3952	82	*	*	*	*	*	*	
3953	102	74.5	15.7	4.9	1.0	2.0	2.0	
3954	222	85.1	5.9	3.6	0.0	2.3	3.2	
4001	1,199	26.3	58.2	13.3	0.0	1.4	0.8	
4002	931	53.5	29.3	13.2	0.6	0.9	2.5	
4103	48	*	*	*	*	*	*	
4106	1,881	8.6	49.3	39.3	0.2	0.9	1.6	
4107	3,675	57.5	26.7	10.7	0.7	2.0	2.3	
4109	1,171	92.1	0.9	3.9	0.2	0.3	2.6	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 91  
Run Date: 10 / 11 / 07

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan Native %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other/Unknown %</b>	
4110	1,949	36.8	7.8	52.5	0.8	0.5	1.6	
4111	66	*	*	*	*	*	*	
4112	944	1.2	0.5	97.9	0.0	0.0	0.4	
4113	915	8.2	16.5	73.0	0.2	1.4	0.7	
4120	2,287	67.6	9.4	18.1	0.3	2.7	2.0	
4160	2,275	46.4	30.2	11.8	0.3	8.5	2.8	
4351	790	93.2	1.1	3.8	0.3	0.5	1.1	
4352	259	83.0	12.0	1.2	0.8	0.0	3.1	
4353	621	76.8	0.8	20.5	0.5	0.5	1.0	
4354	965	90.3	5.3	1.7	0.3	0.8	1.7	
4355	417	89.7	0.2	3.4	2.2	0.7	3.8	
4356	646	85.9	0.5	10.2	0.6	0.9	1.9	
4357	23	*	*	*	*	*	*	
4358	333	2.7	0.0	96.4	0.0	0.0	0.9	
4359	400	66.8	0.5	31.3	0.0	0.5	1.0	
4360	240	94.6	0.4	3.8	0.0	0.4	0.8	
4361	291	93.8	2.4	1.4	0.0	0.7	1.7	
4362	440	96.6	0.0	1.6	0.0	0.0	1.8	
4363	999	94.6	0.8	1.0	0.6	0.4	2.6	
4401	1,407	93.2	0.4	3.1	0.3	0.4	2.7	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 92  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
4501	1,163	97.7	0.2	1.2	0.0	0.2	0.8
4601	1,765	76.0	2.0	19.7	0.1	0.4	1.8
4701	1,212	94.9	0.4	2.1	0.1	1.4	1.1
5050	1,167	78.4	17.1	1.5	0.1	1.5	1.4
5051	3,053	74.3	15.4	6.4	0.0	2.3	1.5
5052	1,854	75.6	15.2	1.7	0.1	5.7	1.7
5054	245	44.9	3.3	50.2	0.0	0.8	0.8
5202	382	94.0	0.3	1.3	0.8	0.8	2.9
5203	213	92.0	0.9	0.9	2.8	0.5	2.8
5204	175	90.9	1.7	0.6	1.7	0.0	5.1
5205	271	94.5	0.0	0.4	2.2	0.4	2.6
5601	1,232	97.5	0.8	0.6	0.1	0.2	0.8
5801	2,340	89.2	3.8	5.0	0.1	0.4	1.5
5803	17	*	*	*	*	*	*
6103	4,923	61.6	28.1	7.0	0.6	0.7	2.0
6301	5,274	45.5	26.9	23.5	0.1	2.5	1.5
6312	3,046	44.5	49.4	1.4	0.1	2.6	2.1
6316	1,212	80.1	8.8	7.3	0.0	1.1	2.6
6320	287	68.6	23.3	2.4	0.7	2.4	2.4
6325	177	89.3	3.4	4.0	0.0	2.3	1.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 93  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
6330	329	28.6	53.8	13.1	0.3	2.1	2.1
6335	692	79.2	13.4	2.3	0.6	3.0	1.4
6340	616	56.8	14.6	21.8	0.3	3.7	2.8
7304	321	93.5	0.6	3.7	0.0	0.0	2.2
7306	3,289	53.4	34.0	11.3	0.0	0.4	1.0
7308	229	92.1	1.3	5.7	0.0	0.0	0.9
7401	2,499	84.8	5.8	5.0	0.8	0.8	2.8
7402	353	92.4	0.8	2.5	0.8	0.3	3.1
7403	97	*	*	*	*	*	*
7404	147	95.9	0.7	2.0	0.0	0.7	0.7
7601	830	95.3	0.5	2.8	0.0	0.1	1.3
7850	1,245	95.6	0.4	2.4	0.0	0.2	1.4
7851	239	97.5	0.0	1.3	0.0	0.4	0.8
7852	131	93.9	1.5	1.5	0.8	0.8	1.5
8150	114	66.7	24.6	3.5	0.0	1.8	3.5
8151	2,985	41.3	44.0	10.4	0.3	1.9	2.1
8152	572	41.4	48.4	5.6	0.0	0.5	4.0
8153	35	*	*	*	*	*	*
8154	441	29.7	14.3	10.2	0.2	45.4	0.2
8155	81	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 94  
Run Date: 10 / 11 / 07

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %	
8156	48	*	*	*	*	*	*	
8157	9	*	*	*	*	*	*	
8201	2,452	64.7	28.1	3.2	0.4	1.8	1.9	
8202	3,302	91.2	3.8	2.6	0.2	1.1	1.2	
8253	606	68.2	26.4	1.0	0.0	0.5	4.0	
8254	685	66.1	20.3	11.2	0.4	0.1	1.8	
8255	3,204	77.3	19.5	1.3	0.0	0.6	1.2	
8257	539	44.7	25.8	0.6	0.2	26.9	1.9	
8258	1,747	73.0	21.4	3.0	0.1	0.5	1.9	
8901	1,252	1.0	96.5	0.6	0.2	0.4	1.3	
8902	634	2.5	95.9	0.6	0.0	0.2	0.8	
8903	1,321	3.0	93.9	1.0	0.1	0.5	1.6	
8904	877	13.5	64.8	2.5	0.6	17.1	1.6	
8905	2,292	22.2	32.4	35.1	0.4	7.2	2.7	
9001	77	*	*	*	*	*	*	
9002	29	*	*	*	*	*	*	
9004	574	0.2	98.3	0.2	0.0	0.5	0.9	
9005	11	*	*	*	*	*	*	
9006	33	*	*	*	*	*	*	
9008	6	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 95  
Run Date: 10 / 11 / 07

<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>						
		<b>White, Not Hispanic</b> %	<b>Black, Not Hispanic</b> %	<b>Hispanic</b> %	<b>American Indian/Alaskan Native</b> %	<b>Asian/Pacific Islander</b> %	<b>All Other/Unknown</b> %	
9009	34	*	*	*	*	*	*	*
9010	280	1.1	97.9	0.4	0.4	0.0	0.4	
9011	14	*	*	*	*	*	*	*
9012	31	*	*	*	*	*	*	*
9014	26	*	*	*	*	*	*	*
9018	10	*	*	*	*	*	*	*
9020	20	*	*	*	*	*	*	*
9022	12	*	*	*	*	*	*	*
9023	65	*	*	*	*	*	*	*
9024	32	*	*	*	*	*	*	*
9035	51	*	*	*	*	*	*	*
9039	47	*	*	*	*	*	*	*
9040	465	1.7	97.0	0.2	0.0	0.0	1.1	
9045	1,507	7.2	53.2	38.2	0.1	0.2	1.1	
9050	2,236	12.7	50.8	31.4	0.2	4.2	0.7	
9051	2,977	11.5	6.6	80.5	0.2	0.2	1.0	
9053	3,136	1.0	96.5	0.5	0.1	0.6	1.2	
9054	819	2.9	91.3	1.8	0.1	1.6	2.2	
9058	558	81.9	15.1	0.2	0.0	2.0	0.9	
9066	3,025	1.4	96.3	0.6	0.0	0.1	1.6	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

Page: 96  
Run Date: 10 / 11 / 07

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaskan Native</b>	<b>Asian/Pacific Islander</b>	<b>All Other/Unknown</b>	
9068	1,724	3.3	94.2	0.3	0.1	0.8	1.3	
9069	2,710	2.5	95.8	0.3	0.0	0.1	1.3	
9070	2,590	3.0	86.8	0.3	0.3	8.4	1.2	
9072	1,861	3.0	90.9	0.2	0.0	4.5	1.4	
9074	1,590	1.0	93.6	0.8	0.1	0.1	4.5	
9079	1,390	0.7	96.8	0.1	0.1	0.4	1.9	
9080	2,313	31.6	64.1	2.8	0.0	0.1	1.3	
9151	1	*	*	*	*	*	*	
Unknown Clinics	468	22.9	53.2	11.8	0.2	0.6	11.3	
<b>Michigan</b>	<b>190,695</b>	<b>56.9</b>	<b>28.3</b>	<b>11.0</b>	<b>0.5</b>	<b>1.7</b>	<b>1.7</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 97  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0101	157	0.0	5.1	15.9	60.5	15.9	2.5	0.0	
0102	499	0.0	5.8	13.6	64.5	13.6	2.4	0.0	
0103	531	0.0	7.5	12.6	66.9	12.2	0.8	0.0	
0104	169	0.0	7.7	16.0	58.0	17.2	1.2	0.0	
0250	199	0.0	9.0	13.6	64.8	12.1	0.5	0.0	
0251	150	0.7	1.3	14.7	69.3	12.7	1.3	0.0	
0252	159	0.0	2.5	11.9	63.5	20.8	1.3	0.0	
0253	182	0.0	1.6	18.1	66.5	12.6	1.1	0.0	
0401	585	0.0	7.0	12.8	66.0	13.3	0.9	0.0	
0402	553	0.2	6.1	12.8	61.8	18.8	0.2	0.0	
0403	157	0.0	6.4	13.4	61.1	17.8	1.3	0.0	
0404	178	0.0	5.1	10.1	66.9	16.3	1.7	0.0	
0501	609	0.3	7.9	13.5	63.2	14.3	0.8	0.0	
0502	3	*	*	*	*	*	*	*	
0503	182	0.0	7.1	14.3	56.0	21.4	1.1	0.0	
0504	701	0.0	3.7	10.7	64.6	19.1	1.9	0.0	
0505	508	0.0	5.3	13.6	66.7	13.4	1.0	0.0	
0506	229	0.4	9.2	11.8	68.6	10.0	0.0	0.0	
0650	1,179	0.1	5.1	13.2	67.4	13.3	0.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 98  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0651	745	0.5	8.7	16.5	62.3	11.3	0.7	0.0	
0652	447	0.2	6.0	14.8	63.1	14.8	1.1	0.0	
0653	479	0.2	5.2	16.1	63.0	14.6	0.8	0.0	
0654	416	0.5	5.3	13.7	65.6	13.9	1.0	0.0	
0655	481	0.6	4.8	13.3	66.3	13.5	1.5	0.0	
0657	126	0.8	4.8	12.7	65.1	16.7	0.0	0.0	
0701	511	0.0	3.9	9.2	63.0	22.1	1.8	0.0	
0702	126	0.0	11.1	14.3	53.2	19.8	1.6	0.0	
0703	276	0.4	5.1	15.6	60.9	16.7	1.4	0.0	
0704	96	*	*	*	*	*	*	*	
0705	189	0.0	5.3	11.1	57.1	24.9	1.6	0.0	
0850	1,247	0.1	4.6	14.5	65.4	14.8	0.6	0.0	
0851	960	0.1	7.1	16.7	62.1	13.3	0.7	0.0	
0901	2,158	0.4	6.9	13.3	62.2	15.6	1.6	0.0	
0902	84	*	*	*	*	*	*	*	
1051	1,626	0.1	4.6	11.7	63.5	19.0	1.0	0.0	
1059	4	*	*	*	*	*	*	*	
1060	1	*	*	*	*	*	*	*	
1101	2,148	0.7	8.4	15.1	60.8	14.2	0.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 99  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1102	1,020	0.6	8.5	15.0	61.5	13.7	0.7	0.0	
1106	120	0.0	7.5	10.0	67.5	13.3	1.7	0.0	
1250	1,139	0.4	8.3	16.9	61.1	12.2	1.1	0.0	
1251	916	0.4	6.1	14.7	65.8	12.3	0.5	0.0	
1252	945	0.2	6.6	12.8	65.0	14.7	0.7	0.0	
1253	879	0.8	8.2	18.0	60.1	12.6	0.3	0.0	
1350	2,870	0.5	8.8	15.9	61.6	12.5	0.7	0.0	
1351	212	0.0	5.2	11.3	68.9	13.7	0.9	0.0	
1352	551	0.2	9.6	13.8	63.7	12.0	0.7	0.0	
1401	325	0.0	4.0	12.9	62.2	18.8	2.2	0.0	
1402	193	0.0	5.2	13.0	58.5	19.7	3.6	0.0	
1403	30	*	*	*	*	*	*	*	*
1404	22	*	*	*	*	*	*	*	*
1405	1	*	*	*	*	*	*	*	*
1501	1,369	0.4	7.4	14.0	62.4	14.6	1.3	0.0	
1502	118	0.0	9.3	19.5	61.9	9.3	0.0	0.0	
1503	1,359	0.3	7.9	13.5	59.1	18.0	1.1	0.0	
1504	2,882	0.6	8.2	13.2	60.0	17.3	0.7	0.0	
1505	700	0.1	5.1	12.1	60.6	20.7	1.3	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 100  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1506	564	0.2	6.0	13.3	61.7	17.4	1.4	0.0	
1507	707	0.3	4.7	13.0	66.5	14.4	1.0	0.1	
1508	303	0.3	8.3	10.2	57.4	22.1	1.7	0.0	
1509	733	0.4	8.2	12.3	62.5	15.4	1.2	0.0	
1510	264	0.4	8.7	19.7	57.2	12.9	1.1	0.0	
1511	598	0.7	5.0	12.9	66.1	15.1	0.3	0.0	
1512	332	0.3	5.4	17.8	59.6	14.8	1.8	0.3	
1701	773	0.1	4.8	10.9	66.4	16.7	1.2	0.0	
1712	138	0.0	6.5	10.1	63.0	18.8	1.4	0.0	
1950	786	0.4	7.3	17.3	62.5	11.6	1.0	0.0	
1952	186	0.5	9.7	18.8	58.6	9.7	2.7	0.0	
1953	445	0.2	3.8	13.0	71.2	11.0	0.7	0.0	
1954	446	0.9	6.7	13.2	71.1	7.8	0.2	0.0	
1955	230	0.0	9.6	13.5	70.0	5.7	1.3	0.0	
1956	14	*	*	*	*	*	*	*	
1957	417	0.0	4.3	15.6	65.2	14.6	0.2	0.0	
1959	229	0.0	6.6	11.8	62.4	19.2	0.0	0.0	
1963	256	0.4	6.3	16.4	65.6	11.3	0.0	0.0	
2150	830	0.1	6.0	11.8	64.2	16.0	1.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 101  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
2151	361	0.0	6.1	16.3	61.5	13.9	2.2	0.0	
2251	606	0.2	5.4	12.4	62.5	18.3	1.2	0.0	
2252	228	0.0	7.0	11.0	65.4	16.2	0.4	0.0	
2551	4,419	1.1	9.2	14.7	59.2	14.6	1.2	0.0	
2552	1,104	1.3	9.5	15.9	58.2	14.2	1.0	0.0	
2554	124	0.0	8.1	18.5	62.9	9.7	0.8	0.0	
2555	4,504	0.3	5.6	12.2	64.8	16.3	0.9	0.0	
3001	1,256	1.8	13.3	15.1	53.8	15.4	0.6	0.0	
3002	238	0.8	8.4	16.4	54.6	18.1	1.7	0.0	
3003	34	*	*	*	*	*	*	*	*
3004	266	0.4	8.6	12.0	57.5	20.3	1.1	0.0	
3007	35	*	*	*	*	*	*	*	*
3009	10	*	*	*	*	*	*	*	*
3010	1	*	*	*	*	*	*	*	*
3201	686	0.0	4.5	10.5	59.2	24.2	1.6	0.0	
3202	23	*	*	*	*	*	*	*	*
3301	4,570	0.3	7.3	13.7	61.2	16.7	0.8	0.0	
3302	476	0.6	7.4	15.5	56.1	19.1	1.3	0.0	
3303	88	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 102  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
3304	213	0.0	0.0	4.2	47.9	47.9	0.0	0.0	
3305	117	0.0	0.0	8.5	65.0	23.9	2.6	0.0	
3306	120	0.0	7.5	12.5	70.8	8.3	0.8	0.0	
3308	72	*	*	*	*	*	*	*	*
3309	49	*	*	*	*	*	*	*	*
3311	21	*	*	*	*	*	*	*	*
3550	105	0.0	2.9	13.3	73.3	9.5	1.0	0.0	
3551	33	*	*	*	*	*	*	*	*
3552	52	*	*	*	*	*	*	*	*
3650	37	*	*	*	*	*	*	*	*
3701	822	0.2	8.8	13.7	63.0	13.3	1.0	0.0	
3702	273	0.0	8.1	17.2	63.4	10.6	0.7	0.0	
3703	126	0.0	4.8	17.5	64.3	12.7	0.8	0.0	
3704	112	0.0	6.3	17.0	67.0	8.9	0.9	0.0	
3805	3,714	0.2	9.5	16.1	61.0	12.5	0.7	0.0	
3950	2,327	0.2	6.1	11.9	63.6	17.1	1.1	0.0	
3952	82	*	*	*	*	*	*	*	*
3953	102	0.0	2.9	12.7	62.7	20.6	1.0	0.0	
3954	222	0.0	5.0	13.1	64.4	17.6	0.0	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 103  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4001	1,199	1.2	11.9	15.1	57.5	13.6	0.7	0.0	
4002	931	0.4	7.9	15.7	61.4	13.3	1.2	0.0	
4103	48	*	*	*	*	*	*	*	*
4106	1,881	1.0	10.2	13.2	58.2	15.6	1.9	0.0	
4107	3,675	0.5	6.8	12.8	62.0	16.9	1.1	0.0	
4109	1,171	0.2	6.7	14.1	60.3	17.2	1.5	0.0	
4110	1,949	0.6	7.9	13.3	60.8	16.4	1.1	0.0	
4111	66	*	*	*	*	*	*	*	*
4112	944	0.1	4.4	9.3	56.7	27.3	2.1	0.0	
4113	915	1.0	8.1	14.2	59.2	16.9	0.5	0.0	
4120	2,287	0.3	7.0	13.9	60.8	16.7	1.4	0.0	
4160	2,275	0.4	5.9	11.7	62.8	18.2	1.0	0.0	
4351	790	0.4	6.6	18.9	62.0	11.3	0.9	0.0	
4352	259	0.8	7.7	17.0	61.4	13.1	0.0	0.0	
4353	621	0.5	9.0	16.6	61.7	11.6	0.6	0.0	
4354	965	0.1	8.1	15.0	61.8	14.2	0.8	0.0	
4355	417	0.2	6.5	12.2	64.0	15.3	1.7	0.0	
4356	646	0.0	4.5	13.5	64.9	15.8	1.4	0.0	
4357	23	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 104  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4358	333	1.5	7.2	10.8	58.3	21.0	1.2	0.0	
4359	400	0.3	7.3	16.0	59.3	16.3	1.0	0.0	
4360	240	0.4	7.5	13.8	63.3	14.6	0.4	0.0	
4361	291	1.0	10.7	13.7	61.2	13.4	0.0	0.0	
4362	440	0.2	8.4	14.5	64.3	11.8	0.7	0.0	
4363	999	0.3	7.1	18.1	63.3	10.5	0.7	0.0	
4401	1,407	0.0	4.3	15.1	63.2	16.5	0.9	0.0	
4501	1,163	0.4	5.2	12.4	65.0	16.5	0.5	0.0	
4601	1,765	0.1	8.3	14.7	63.5	12.6	0.8	0.0	
4701	1,212	0.0	3.3	16.0	62.0	18.0	0.7	0.0	
5050	1,167	0.3	4.1	10.7	59.4	23.7	1.7	0.0	
5051	3,053	0.1	4.6	12.5	61.8	19.4	1.7	0.0	
5052	1,854	0.2	3.7	10.0	59.1	25.2	1.8	0.0	
5054	245	0.0	7.3	10.6	61.6	19.6	0.8	0.0	
5202	382	0.0	1.6	8.9	67.8	20.9	0.8	0.0	
5203	213	0.0	4.7	10.3	67.6	16.0	1.4	0.0	
5204	175	0.0	4.6	10.3	69.7	14.3	1.1	0.0	
5205	271	0.4	1.5	10.3	64.2	21.0	1.5	1.1	
5601	1,232	0.2	5.7	14.0	64.5	14.9	0.6	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 105  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
5801	2,340	0.3	6.7	15.4	62.8	14.1	0.6	0.0	
5803	17	*	*	*	*	*	*	*	*
6103	4,923	0.4	8.5	14.1	63.1	13.1	0.8	0.0	
6301	5,274	0.4	7.3	13.0	60.3	17.8	1.3	0.0	
6312	3,046	0.2	3.8	8.9	57.4	27.1	2.6	0.0	
6316	1,212	0.2	3.5	9.9	66.3	19.1	1.0	0.0	
6320	287	0.0	3.5	6.6	61.7	25.4	2.8	0.0	
6325	177	0.0	4.0	14.7	63.3	15.3	2.8	0.0	
6330	329	0.3	8.8	18.8	54.4	14.6	3.0	0.0	
6335	692	0.3	3.3	9.7	59.7	24.1	2.9	0.0	
6340	616	0.3	4.1	14.6	64.6	15.7	0.6	0.0	
7304	321	0.0	4.0	13.7	59.5	19.9	2.8	0.0	
7306	3,289	0.3	6.9	14.0	61.1	16.4	1.3	0.0	
7308	229	0.0	4.4	6.1	62.4	26.2	0.9	0.0	
7401	2,499	0.2	7.1	13.7	64.2	14.1	0.6	0.0	
7402	353	0.3	5.9	14.4	57.5	20.7	1.1	0.0	
7403	97	*	*	*	*	*	*	*	*
7404	147	0.0	6.1	14.3	59.9	18.4	1.4	0.0	
7601	830	0.0	7.1	11.7	64.8	15.2	1.2	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 106  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
7850	1,245	0.1	5.7	16.5	63.3	12.9	1.5	0.0	
7851	239	0.0	3.8	13.8	65.3	15.9	1.3	0.0	
7852	131	1.5	3.1	8.4	69.5	15.3	2.3	0.0	
8150	114	0.0	0.9	4.4	61.4	29.8	2.6	0.9	
8151	2,985	0.1	2.1	7.7	67.2	21.7	1.2	0.0	
8152	572	2.3	25.3	40.4	32.0	0.0	0.0	0.0	
8153	35	*	*	*	*	*	*	*	*
8154	441	0.0	2.0	2.7	57.4	36.7	1.1	0.0	
8155	81	*	*	*	*	*	*	*	*
8156	48	*	*	*	*	*	*	*	*
8157	9	*	*	*	*	*	*	*	*
8201	2,452	0.3	6.4	13.3	59.1	20.1	0.8	0.0	
8202	3,302	0.1	2.5	8.6	60.2	25.7	2.9	0.0	
8253	606	0.3	4.8	11.2	64.7	17.8	1.2	0.0	
8254	685	0.1	7.2	13.7	61.6	16.6	0.7	0.0	
8255	3,204	0.3	6.7	13.1	63.0	15.9	1.1	0.0	
8257	539	0.2	4.3	10.4	58.4	23.2	3.5	0.0	
8258	1,747	0.2	7.0	11.5	61.8	18.5	1.1	0.0	
8901	1,252	1.8	12.1	12.8	55.7	16.6	1.0	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 107  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
8902	634	1.4	8.0	12.9	53.9	22.6	1.1	0.0	
8903	1,321	1.1	11.4	14.7	50.9	20.7	1.2	0.0	
8904	877	1.0	5.6	10.3	58.0	23.8	1.3	0.0	
8905	2,292	0.7	7.5	11.3	54.8	24.0	1.7	0.0	
9001	77	*	*	*	*	*	*	*	*
9002	29	*	*	*	*	*	*	*	*
9004	574	1.0	9.2	11.7	54.5	22.5	1.0	0.0	
9005	11	*	*	*	*	*	*	*	*
9006	33	*	*	*	*	*	*	*	*
9008	6	*	*	*	*	*	*	*	*
9009	34	*	*	*	*	*	*	*	*
9010	280	1.1	8.2	14.6	58.6	16.4	0.7	0.4	
9011	14	*	*	*	*	*	*	*	*
9012	31	*	*	*	*	*	*	*	*
9014	26	*	*	*	*	*	*	*	*
9018	10	*	*	*	*	*	*	*	*
9020	20	*	*	*	*	*	*	*	*
9022	12	*	*	*	*	*	*	*	*
9023	65	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 108  
Run Date: 10 / 11 / 07

Clinic	Number Included in Analysis	Comparison of Age Distribution <sup>(2)</sup>							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
9024	32	*	*	*	*	*	*	*	*
9035	51	*	*	*	*	*	*	*	*
9039	47	*	*	*	*	*	*	*	*
9040	465	1.7	12.7	16.1	50.8	17.4	1.3	0.0	
9045	1,507	3.7	17.0	13.2	46.4	18.2	1.3	0.1	
9050	2,236	0.8	8.0	10.8	56.9	21.4	2.1	0.0	
9051	2,977	0.3	6.7	10.5	60.0	21.0	1.4	0.0	
9053	3,136	0.4	8.5	13.8	54.4	21.3	1.5	0.0	
9054	819	0.9	7.1	10.3	57.9	22.5	1.3	0.1	
9058	558	0.2	2.7	5.4	57.3	29.9	4.5	0.0	
9066	3,025	0.6	9.1	15.2	53.6	20.1	1.4	0.0	
9068	1,724	0.6	7.7	13.9	59.4	17.2	1.2	0.0	
9069	2,710	0.6	7.6	13.5	56.6	20.4	1.3	0.0	
9070	2,590	0.7	9.2	12.7	56.3	19.8	1.4	0.0	
9072	1,861	1.0	10.4	15.0	53.0	19.1	1.5	0.0	
9074	1,590	0.6	7.6	12.8	57.7	19.7	1.4	0.0	
9079	1,390	0.6	5.8	10.3	54.6	26.8	2.0	0.0	
9080	2,313	0.4	5.4	10.7	57.0	24.8	1.6	0.0	
9151	1	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Page: 109  
Run Date: 10 / 11 / 07

<b>Comparison of Age Distribution</b> <sup>(2)</sup>									
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>	
Unknown Clinics	468	1.1	9.4	13.2	54.1	20.3	1.9	0.0	
<b>Michigan</b>	<b>190,695</b>	<b>0.4</b>	<b>7.0</b>	<b>13.2</b>	<b>60.6</b>	<b>17.5</b>	<b>1.2</b>	<b>0.0</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 110  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	150	7.3 (12)	52.0 (161)	129	29.5 (113)	49.6 (149)	16	* * *	41	* * *
0102	472	8.9 (34)	50.0 (141)	398	22.9 (46)	52.3 (161)	76	* * *	95	* * *
0103	509	13.6 (152)	47.5 (117)	445	21.1 (21)	49.0 (139)	67	* * *	116	37.1 (55)
0104	161	18.6 (178)	37.9 (14)	142	21.8 (35)	46.5 (97)	14	* * *	31	* * *
0250	189	11.1 (92)	45.0 (85)	160	42.5 (153)	30.6 (17)	15	* * *	45	* * *
0251	142	11.3 (98)	50.0 (141)	123	39.8 (148)	42.3 (49)	19	* * *	33	* * *
0252	130	6.2 (5)	49.2 (133)	109	45.0 (156)	36.7 (25)	12	* * *	44	* * *
0253	176	8.0 (20)	43.8 (62)	151	66.2 (170)	14.6 (3)	17	* * *	39	* * *
0401	566	12.4 (121)	46.5 (102)	495	25.3 (76)	45.5 (82)	66	* * *	192	24.0 (16)
0402	538	10.2 (69)	47.8 (121)	464	25.6 (83)	45.7 (88)	64	* * *	87	* * *
0403	151	15.9 (174)	43.7 (60)	139	27.3 (98)	42.4 (53)	14	* * *	33	* * *
0404	171	9.9 (61)	55.0 (177)	153	27.5 (101)	45.1 (77)	22	* * *	40	* * *
0501	582	12.7 (134)	41.9 (37)	535	19.6 (7)	47.9 (120)	92	* * *	159	57.2 (98)
0502	3	* * * * *	*	3	* * * * *	*	1	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 111  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
0503	161	14.3 (164)	36.6 (7)	149	32.2 (132)	39.6 (31)	24	* * *	42	* * *
0504	664	10.1 (66)	39.8 (22)	595	30.4 (121)	40.3 (35)	79	* * *	177	57.6(100)
0505	480	11.0 (90)	45.6 (93)	434	54.6 (164)	25.3 (8)	65	* * *	137	67.9(102)
0506	216	12.5(124)	47.2(112)	186	47.3 (159)	29.6 (15)	30	* * *	44	* * *
0650	1,066	9.8 (58)	49.1 (131)	967	25.5 (79)	47.2 (110)	172	21.5 (16)	319	23.8 (15)
0651	699	10.9 (85)	47.4 (116)	604	24.8 (70)	47.5(116)	96	* * *	249	28.9 (33)
0652	412	10.0 (63)	48.1 (125)	352	30.4 (121)	45.2 (79)	66	* * *	120	38.3 (62)
0653	454	10.1 (66)	46.5 (102)	409	28.9 (109)	42.8 (57)	78	* * *	89	* * *
0654	396	12.4(121)	49.5(137)	308	27.6 (103)	49.7 (151)	53	* * *	94	* * *
0655	454	12.8(137)	44.1 (70)	411	21.7 (31)	47.2 (110)	78	* * *	171	22.8 (12)
0657	123	11.4(101)	39.0 (19)	110	25.5 (79)	45.5 (82)	16	* * *	57	* * *
0701	469	19.0(179)	30.3 (4)	380	50.0 (162)	27.1 (9)	64	* * *	165	50.3 (88)
0702	118	14.4(167)	44.9 (84)	99	* * * * *	* * *	12	* * *	24	* * *
0703	247	13.0(139)	40.5 (25)	225	20.9 (18)	50.2 (154)	23	* * *	64	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 112  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>										
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0704	91	*	*	79	*	*	10	*	20	*	*
0705	176	12.5 (124)	42.0 (40)	158	61.4 (169)	20.9 (4)	21	*	53	*	*
0850	1,171	11.1 (92)	46.5 (102)	987	29.7 (115)	43.3 (64)	173	19.1 (9)	342	36.0 (54)	
0851	913	11.3 (98)	41.6 (35)	809	34.4 (138)	41.4 (43)	110	25.5 (29)	254	41.3 (71)	
0901	1,983	12.6 (129)	46.7 (107)	1,731	26.2 (95)	45.6 (85)	292	15.8 (7)	509	18.5 (6)	
0902	79	*	*	67	*	*	12	*	19	*	*
1051	1,532	11.0 (90)	41.6 (35)	1,332	21.2 (23)	47.4 (114)	206	32.0 (48)	510	33.7 (48)	
1059	4	*	*	4	*	*	2	*	3	*	*
1060	1	*	*	1	*	*	0	*	0	*	*
1101	2,065	11.2 (94)	50.3 (148)	1,723	25.2 (74)	47.4 (114)	414	41.5 (71)	658	42.9 (75)	
1102	947	14.0 (158)	42.9 (49)	799	22.3 (41)	52.2 (160)	177	20.9 (12)	307	54.1 (96)	
1106	112	12.5 (124)	35.7 (6)	96	*	*	29	*	41	*	*
1250	1,048	14.5 (169)	41.3 (33)	897	30.4 (121)	43.5 (67)	156	24.4 (22)	232	35.3 (53)	
1251	868	13.2 (143)	43.8 (62)	781	23.0 (47)	49.0 (139)	179	26.8 (32)	267	32.2 (42)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 113  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
1252	889	10.8 (84)	49.0 (130)	763	27.7 (104)	45.6 (85)	161	44.1 (77)	210	41.0 (70)
1253	832	12.6 (129)	43.3 (52)	711	45.6 (157)	34.7 (22)	142	28.9 (40)	181	38.1 (60)
1350	2,717	10.6 (78)	44.7 (78)	2,395	22.5 (43)	49.0 (139)	516	31.8 (47)	536	50.9 (90)
1351	200	9.0 (37)	40.5 (25)	179	23.5 (55)	53.6 (166)	44	* * *	30	* * *
1352	536	8.8 (31)	45.0 (85)	450	26.0 (92)	46.0 (90)	83	* * *	95	* * *
1401	309	9.4 (45)	49.2 (133)	260	22.3 (41)	48.1 (123)	46	* * *	80	* * *
1402	173	6.4 (7)	53.2 (173)	156	25.6 (83)	44.9 (76)	27	* * *	41	* * *
1403	29	* * * * *		26	* * * * *		4	* * *	8	* * *
1404	16	* * * * *		8	* * * * *		2	* * *	1	* * *
1405	1	* * * * *		0	* * * * *		0	* * *	0	* * *
1501	1,244	9.7 (54)	53.0 (172)	1,054	30.1 (119)	40.4 (37)	182	33.5 (56)	220	52.7 (94)
1502	113	12.4 (121)	40.7 (28)	81	* * * * *		21	* * *	20	* * *
1503	1,156	9.5 (48)	47.1 (109)	935	57.2 (166)	27.3 (10)	184	25.0 (27)	230	43.0 (76)
1504	2,692	10.9 (85)	42.6 (44)	2,333	51.4 (163)	27.9 (11)	473	30.2 (43)	703	30.3 (38)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 114  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
1505	643	13.7 (155)	37.9 (14)	579	27.3 (98)	44.7 (74)	127	21.3 (15)	164	34.8 (51)
1506	519	13.3 (147)	41.2 (32)	440	20.0 (11)	49.3 (144)	75	* * *	78	* * *
1507	675	9.8 (58)	43.7 (60)	593	19.6 (7)	47.9 (120)	91	* * *	155	40.0 (67)
1508	193	7.3 (12)	48.2 (126)	130	47.7 (160)	23.1 (6)	27	* * *	45	* * *
1509	669	6.6 (9)	52.9 (170)	559	30.1 (119)	41.1 (41)	84	* * *	102	39.2 (64)
1510	245	17.1 (177)	39.2 (20)	206	23.3 (54)	48.1 (123)	37	* * *	59	* * *
1511	562	9.4 (45)	50.2 (146)	496	21.0 (20)	53.4 (165)	76	* * *	105	44.8 (79)
1512	312	14.1 (159)	45.8 (96)	279	21.5 (28)	47.0 (108)	55	* * *	71	* * *
1701	751	8.8 (31)	49.3 (136)	652	20.7 (15)	50.5 (155)	98	* * *	99	* * *
1712	130	6.2 (5)	50.8 (151)	117	25.6 (83)	53.8 (167)	14	* * *	17	* * *
1950	762	14.2 (162)	45.3 (91)	662	24.5 (65)	46.5 (97)	122	13.1 (3)	215	14.0 (2)
1952	181	12.7 (134)	45.9 (97)	168	26.8 (97)	42.9 (59)	40	* * *	50	* * *
1953	432	14.1 (159)	43.8 (62)	399	24.6 (67)	46.9 (105)	70	* * *	116	19.8 (9)
1954	421	9.7 (54)	52.7 (169)	362	25.7 (88)	47.5 (116)	48	* * *	80	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 115  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
1955	215	13.0 (139)	40.5 (25)	185	25.9 (90)	48.1 (123)	34	* * *	44	* * *
1956	11	* * * *	* *	8	* * * *	* *	0	* * *	3	* * *
1957	384	13.5 (149)	41.1 (31)	348	20.7 (15)	48.3 (126)	64	* * *	86	* * *
1959	206	8.7 (30)	51.0 (153)	180	22.8 (44)	52.8 (163)	40	* * *	56	* * *
1963	234	6.4 (7)	49.6 (139)	199	25.1 (73)	46.7 (102)	36	* * *	50	* * *
2150	788	9.6 (51)	45.1 (89)	667	20.2 (13)	53.1 (164)	114	12.3 (2)	183	28.4 (30)
2151	329	10.9 (85)	42.2 (42)	288	23.6 (58)	48.6 (132)	48	* * *	65	* * *
2251	553	11.6 (107)	44.8 (80)	484	19.6 (7)	49.4 (146)	58	* * *	167	19.2 (7)
2252	211	9.5 (48)	43.6 (55)	183	41.5 (152)	33.9 (20)	17	* * *	51	* * *
2551	4,157	11.8 (112)	46.8 (108)	3,600	23.9 (62)	48.3 (126)	895	47.6 (80)	806	40.0 (67)
2552	1,050	10.9 (85)	47.2 (112)	877	21.7 (31)	48.9 (137)	246	51.6 (82)	152	43.4 (77)
2554	124	16.1 (175)	44.4 (71)	102	59.8 (167)	21.6 (5)	14	* * *	17	* * *
2555	4,202	13.4 (148)	44.4 (71)	3,630	21.8 (35)	48.3 (126)	860	33.4 (55)	873	27.6 (27)
3001	1,182	9.1 (39)	52.2 (165)	972	28.9 (109)	43.3 (64)	163	35.6 (63)	587	30.7 (39)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 116  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
3002	226	9.7 (54)	54.4 (176)	193	29.0 (111)	43.0 (60)	28	* * *	120	49.2 (87)
3003	28	* * * *	* *	18	* * * *	* *	3	* * *	8	* * *
3004	252	5.6 (3)	51.6 (158)	216	23.1 (49)	38.9 (30)	18	* * *	39	* *
3007	29	* * * *	* *	24	* * * *	* *	5	* * *	15	* *
3009	7	* * * *	* *	3	* * * *	* *	0	* * *	0	* *
3010	0	* * * *	* *	0	* * * *	* *	0	* * *	0	* *
3201	664	11.7(110)	47.9(123)	579	25.6 (83)	47.8 (119)	96	* * *	87	* *
3202	21	* * * *	* *	18	* * * *	* *	2	* * *	3	* *
3301	4,196	11.2 (94)	43.6 (55)	3,679	46.4 (158)	32.6 (19)	877	30.2 (43)	1,485	47.3 (82)
3302	434	9.9 (61)	49.8 (140)	388	32.2 (132)	42.3 (49)	81	* * *	163	52.1 (92)
3303	80	* * * *	* *	72	* * * *	* *	20	* * *	27	* *
3304	200	25.5(181)	24.0 (2)	174	39.7 (147)	29.3 (13)	21	* * *	73	* *
3305	106	13.2(143)	34.9 (5)	94	* * * *	* *	20	* * *	33	* *
3306	107	11.2 (94)	52.3 (166)	98	* * * *	* *	15	* * *	48	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 117  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester			Number	% (Rank)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)		Number	% (Rank)	
3308	68	*	*	62	*	*	14	*	*	26	*	*
3309	46	*	*	39	*	*	7	*	*	14	*	*
3311	19	*	*	19	*	*	2	*	*	9	*	*
3550	101	7.9 (19)	60.4 (181)	87	*	*	9	*	*	11	*	*
3551	31	*	*	29	*	*	6	*	*	10	*	*
3552	52	*	*	46	*	*	7	*	*	18	*	*
3650	29	*	*	9	*	*	5	*	*	4	*	*
3701	780	11.8 (112)	44.4 (71)	666	24.6 (67)	47.3 (112)	133	21.1 (14)		175	46.3 (80)	
3702	258	11.6 (107)	45.0 (85)	220	24.1 (63)	42.3 (49)	61	*	*	76	*	*
3703	125	13.6 (152)	47.2 (112)	113	26.5 (96)	46.9 (105)	25	*	*	34	*	*
3704	106	12.3 (120)	49.1 (131)	94	*	*	22	*	*	24	*	*
3805	3,491	12.6 (129)	42.7 (45)	3,126	23.5 (55)	48.3 (126)	505	26.7 (31)		914	51.6 (91)	
3950	2,227	9.8 (58)	47.6 (118)	1,904	23.7 (60)	49.5 (147)	318	42.5 (76)		444	38.1 (60)	
3952	79	*	*	60	*	*	13	*	*	13	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 118  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum		
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
3953	100	10.0 (63)	45.0 (85)	91	*	*	*	*	16	*	*	
3954	215	11.6 (107)	48.8 (128)	190	15.8 (1)	55.8 (168)	30	*	*	43	*	*
4001	1,122	9.1 (39)	50.2 (146)	859	34.7 (139)	41.8 (45)	151	41.7 (73)	157	53.5 (95)		
4002	877	10.7 (82)	47.1 (109)	699	30.6 (125)	43.1 (63)	118	36.4 (65)	133	47.4 (83)		
4103	44	*	*	38	*	*	*	*	6	*	*	
4106	1,762	8.1 (21)	50.9 (152)	1,517	29.9 (117)	40.2 (34)	278	29.5 (41)	335	34.6 (50)		
4107	3,384	11.4 (101)	43.6 (55)	2,882	23.5 (55)	46.3 (95)	539	22.8 (19)	806	21.6 (11)		
4109	1,086	12.7 (134)	44.8 (80)	961	21.9 (38)	47.3 (112)	185	14.6 (5)	279	14.3 (3)		
4110	1,821	8.8 (31)	46.2 (100)	1,584	28.6 (107)	40.5 (39)	203	22.7 (18)	362	27.6 (27)		
4111	59	*	*	52	*	*	*	*	9	*	*	
4112	842	3.6 (1)	51.0 (153)	744	59.9 (168)	13.4 (1)	109	28.4 (36)	293	65.9(101)		
4113	890	6.0 (4)	45.7 (95)	765	30.7 (127)	37.1 (27)	76	*	*	169	46.7 (81)	
4120	2,192	11.7 (110)	42.8 (46)	1,957	21.1 (21)	50.1 (153)	347	21.0 (13)	468	24.4 (17)		
4160	2,137	11.9 (114)	40.2 (23)	1,872	25.5 (79)	43.6 (69)	360	21.7 (17)	460	17.0 (4)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 119  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
4351	762	12.2 (119)	42.8 (46)	668	21.7 (31)	49.1 (142)	112	27.7 (35)	204	27.0 (25)
4352	244	6.6 (9)	52.9 (170)	204	21.6 (29)	49.5 (147)	36	* * *	61	* * *
4353	584	9.6 (51)	47.8 (121)	500	21.2 (23)	48.6 (132)	83	* * *	152	48.7 (85)
4354	920	11.4 (101)	47.1 (109)	796	26.0 (92)	44.7 (74)	118	29.7 (42)	227	33.5 (46)
4355	390	14.6 (170)	42.8 (46)	347	24.5 (65)	55.9 (169)	70	* * *	97	* * *
4356	606	10.6 (78)	47.7 (120)	527	22.2 (39)	49.1 (142)	95	* * *	179	38.0 (59)
4357	11	* * * * *		7	* * * * *		2	* * *	2	* * *
4358	276	8.3 (23)	52.5 (168)	221	39.8 (148)	23.5 (7)	25	* * *	54	* * *
4359	375	10.4 (74)	48.8 (128)	340	25.3 (76)	43.5 (67)	44	* * *	89	* * *
4360	234	11.5 (106)	43.6 (55)	203	23.6 (58)	46.8 (104)	25	* * *	65	* * *
4361	280	15.0 (172)	40.7 (28)	244	20.1 (12)	47.5 (116)	41	* * *	72	* * *
4362	426	11.3 (98)	44.6 (76)	387	20.7 (15)	45.5 (82)	67	* * *	134	29.9 (36)
4363	958	10.5 (77)	44.5 (75)	837	19.6 (7)	50.9 (156)	129	28.7 (39)	249	27.7 (29)
4401	746	13.1 (141)	44.8 (80)	669	23.2 (51)	48.6 (132)	226	25.7 (30)	593	37.9 (58)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 120  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
4501	1,021	14.4 (167)	43.8 (62)	904	23.7 (60)	47.0 (108)	146	15.1 (6)	243	17.3 (5)
4601	1,693	12.5 (124)	43.8 (62)	1,482	23.0 (47)	48.0 (122)	217	20.7 (11)	337	42.1 (73)
4701	1,132	13.8 (156)	38.8 (18)	990	20.9 (18)	52.6 (162)	265	11.3 (1)	426	13.4 (1)
5050	1,119	11.4 (101)	43.9 (67)	319	31.3 (130)	42.6 (54)	283	35.0 (59)	204	29.9 (36)
5051	2,871	13.8 (156)	40.7 (28)	959	30.6 (125)	42.1 (47)	716	33.1 (53)	673	25.0 (20)
5052	1,759	12.5 (124)	41.3 (33)	474	30.0 (118)	40.3 (35)	462	37.2 (67)	340	25.3 (22)
5054	228	9.2 (42)	43.0 (50)	114	33.3 (136)	40.4 (37)	43	* * *	76	* * *
5202	357	8.4 (26)	45.1 (89)	316	21.2 (23)	46.5 (97)	31	* * *	81	* * *
5203	196	9.7 (54)	58.7 (180)	173	18.5 (4)	57.8 (170)	22	* * *	47	* * *
5204	170	9.4 (45)	37.6 (10)	147	19.0 (5)	57.8 (170)	20	* * *	35	* * *
5205	253	12.6 (129)	41.9 (37)	228	23.2 (51)	43.0 (60)	32	* * *	66	* * *
5601	908	13.2 (143)	48.5 (127)	780	21.7 (31)	48.6 (132)	162	23.5 (20)	311	37.6 (57)
5801	2,204	14.1 (159)	42.2 (42)	1,886	23.1 (49)	46.2 (94)	355	24.8 (26)	340	27.1 (26)
5803	17	* * * * *		14	* * * * *		0	* * *	4	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 121  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
6103	4,569	10.3 (72)	48.0 (124)	4,103	22.2 (39)	49.6 (149)	721	40.8 (70)	378	43.9 (78)
6301	4,924	10.1 (66)	46.0 (99)	4,307	25.2 (74)	45.1 (77)	1,056	32.3 (51)	1,071	28.8 (32)
6312	2,838	12.1 (117)	44.6 (76)	2,497	25.4 (78)	45.9 (89)	643	35.1 (60)	785	29.4 (34)
6316	1,140	14.2 (162)	40.2 (23)	1,017	17.5 (2)	51.2 (158)	258	24.4 (22)	368	19.8 (9)
6320	259	13.5 (149)	37.8 (13)	208	25.5 (79)	43.3 (64)	72	* *	70	* *
6325	170	11.2 (94)	44.7 (78)	136	18.4 (3)	46.3 (95)	26	* *	41	* *
6330	309	10.4 (74)	47.6 (118)	226	25.7 (88)	42.0 (46)	85	* *	64	* *
6335	642	12.6 (129)	37.7 (12)	513	21.8 (35)	45.2 (79)	134	28.4 (36)	132	34.1 (49)
6340	568	12.0 (116)	41.9 (37)	444	27.3 (98)	42.6 (54)	144	28.5 (38)	121	25.6 (23)
7304	304	10.2 (69)	53.3 (174)	258	19.4 (6)	51.9 (159)	58	* *	73	* *
7306	3,152	8.2 (22)	51.8 (160)	2,730	20.5 (14)	51.1 (157)	498	31.1 (45)	717	41.8 (72)
7308	219	4.6 (2)	50.7 (150)	197	21.3 (26)	45.2 (79)	36	* *	70	* *
7401	2,390	13.6 (152)	43.9 (67)	2,093	21.6 (29)	48.9 (137)	425	35.3 (61)	1,404	39.4 (65)
7402	338	16.6 (176)	37.9 (14)	298	24.8 (70)	48.3 (126)	84	* *	194	28.4 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 122  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
7403	90	*	*	82	*	*	29	*	53	*	*	
7404	136	6.6 (9)	38.2 (17)	122	21.3 (26)	46.7 (102)	29	*	73	*	*	
7601	796	11.4 (101)	44.0 (69)	725	23.2 (51)	46.6 (100)	150	18.0 (8)	272	23.5 (14)		
7850	529	11.9 (114)	45.9 (97)	472	24.2 (64)	46.0 (90)	150	32.0 (48)	254	32.3 (43)		
7851	192	13.5 (149)	43.2 (51)	166	25.9 (90)	48.8 (136)	30	*	60	*	*	
7852	105	14.3 (164)	44.8 (80)	90	*	*	21	*	33	*	*	
8150	101	10.9 (85)	43.6 (55)	75	*	*	21	*	20	*	*	
8151	2,652	10.3 (72)	46.6 (106)	2,213	29.6 (114)	44.3 (72)	537	41.5 (71)	615	33.2 (45)		
8152	549	13.1 (141)	37.2 (9)	463	33.0 (135)	41.7 (44)	41	*	69	*	*	
8153	32	*	*	29	*	*	9	*	7	*	*	
8154	406	21.7 (180)	19.7 (1)	329	26.1 (94)	29.8 (16)	83	*	142	32.4 (44)		
8155	76	*	*	71	*	*	13	*	15	*	*	
8156	43	*	*	37	*	*	11	*	20	*	*	
8157	8	*	*	6	*	*	2	*	0	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 123  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
8201	2,300	13.2 (143)	42.0 (40)	1,737	24.6 (67)	46.9 (105)	726	24.5 (24)	700	22.9 (13)
8202	3,080	10.6 (78)	37.0 (8)	2,532	40.7 (151)	31.9 (18)	589	31.4 (46)	626	26.8 (24)
8253	541	10.4 (74)	43.3 (52)	441	27.7 (104)	46.0 (90)	122	25.4 (28)	154	24.7 (18)
8254	638	14.3 (164)	46.4 (101)	472	28.4 (106)	46.6 (100)	166	24.7 (25)	166	24.7 (18)
8255	2,946	12.9 (138)	43.5 (54)	2,248	31.0 (128)	42.2 (48)	737	20.4 (10)	832	19.6 (8)
8257	500	14.8 (171)	37.6 (10)	377	40.3 (150)	35.3 (23)	109	32.1 (50)	125	33.6 (47)
8258	1,672	10.6 (78)	45.3 (91)	1,424	22.8 (44)	49.9 (152)	351	34.5 (57)	319	29.5 (35)
8901	1,139	9.1 (39)	52.0 (161)	953	35.4 (142)	39.7 (33)	99	* *	60	* *
8902	592	10.0 (63)	51.4 (157)	500	28.8 (108)	43.0 (60)	36	* *	21	* *
8903	1,224	7.7 (17)	54.2 (175)	1,031	33.9 (137)	42.7 (56)	155	51.0 (81)	78	* *
8904	806	8.6 (28)	50.1 (143)	699	49.2 (161)	29.5 (14)	129	36.4 (65)	134	52.2 (93)
8905	2,064	9.3 (44)	47.3 (115)	1,723	36.3 (144)	37.3 (28)	97	* *	42	* *
9001	75	* *	* *	73	* *	* *	15	* *	1	* *
9002	27	* *	* *	23	* *	* *	4	* *	6	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 124  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
9004	563	8.3 (23)	55.1 (178)	524	31.1 (129)	46.0 (90)	121	34.7 (58)	165	50.3 (88)
9005	11	*	*	11	*	*	6	*	5	*
9006	33	*	*	29	*	*	8	*	4	*
9008	6	*	*	5	*	*	0	*	1	*
9009	33	*	*	29	*	*	12	*	10	*
9010	267	8.6 (28)	51.7 (159)	240	30.4 (121)	44.2 (71)	47	*	42	*
9011	14	*	*	14	*	*	2	*	2	*
9012	29	*	*	28	*	*	6	*	8	*
9014	26	*	*	26	*	*	3	*	6	*
9018	10	*	*	9	*	*	3	*	2	*
9020	20	*	*	19	*	*	2	*	9	*
9022	11	*	*	9	*	*	1	*	0	*
9023	61	*	*	56	*	*	6	*	1	*
9024	31	*	*	28	*	*	6	*	5	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 125  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>																
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester			Number	% (Rank)						
9035	51	*	*	*	*		44	*	*	*	*	10	*	*	13	*	*
9039	43	*	*	*	*		38	*	*	*	*	13	*	*	16	*	*
9040	459	8.9 (34)	49.5 (137)	410	24.9 (72)	49.3 (144)	114	27.2 (33)	92	*	*	92	*	*			
9045	1,378	10.2 (69)	46.5 (102)	1,211	35.5 (143)	37.6 (29)	209	14.4 (4)	166	31.3 (41)							
9050	2,044	9.6 (51)	45.6 (93)	1,776	31.8 (131)	41.0 (40)	285	32.3 (51)	296	39.5 (66)							
9051	2,835	9.5 (48)	39.7 (21)	2,451	34.8 (140)	36.7 (25)	293	23.5 (20)	1,051	25.2 (21)							
9053	3,004	9.0 (37)	51.0 (153)	2,544	55.0 (165)	28.1 (12)	649	39.4 (69)	667	35.2 (52)							
9054	786	8.9 (34)	51.3 (156)	683	29.3 (112)	44.4 (73)	138	44.2 (79)	103	57.3 (99)							
9058	536	15.5 (173)	29.9 (3)	445	68.1 (171)	13.9 (2)	84	*	*	65	*	*					
9066	2,897	10.7 (82)	50.5 (149)	2,501	36.7 (145)	41.3 (42)	519	35.5 (62)	657	55.6 (97)							
9068	1,659	9.2 (42)	52.1 (163)	1,456	37.8 (146)	39.6 (31)	288	42.4 (75)	290	49.0 (86)							
9069	2,528	7.8 (18)	52.1 (163)	2,208	34.9 (141)	42.3 (49)	556	33.1 (53)	983	31.1 (40)							
9070	2,487	12.1 (117)	49.2 (133)	2,098	43.3 (154)	35.5 (24)	398	35.9 (64)	398	38.4 (63)							
9072	1,790	8.4 (26)	50.1 (143)	1,562	32.2 (132)	42.8 (57)	285	42.1 (74)	510	48.2 (84)							

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Page: 126  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Health Indicators <sup>(2, 3)</sup>									
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Postpartum
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
9074	1,545	8.3 (23)	52.4 (167)	1,388	25.6 (83)	48.3 (126)	330	37.9 (68)	285	40.7 (69)
9079	1,347	7.6 (15)	55.7 (179)	1,193	29.7 (115)	45.6 (85)	296	27.4 (34)	265	42.6 (74)
9080	2,239	7.4 (14)	50.1 (143)	1,915	27.5 (101)	43.7 (70)	354	44.1 (77)	270	37.4 (56)
9151	1	*	*	*	*	*	0	*	*	*
Unknown Clinics	381	7.6 (15)	44.4 (71)	285	44.2 (155)	34.4 (21)	91	*	*	*
<b>Michigan</b>	<b>177,236</b>	<b>11.0</b>	<b>45.8</b>	<b>149,207</b>	<b>29.3</b>	<b>43.5</b>	<b>30,973</b>	<b>30.7</b>	<b>42,404</b>	<b>34.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 127  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
0101	141	90.1 (5)	0.0 (1)	152	45.4 (37)	
0102	438	87.7 (13)	3.4 (17)	484	42.4 (56)	
0103	481	87.5 (14)	4.0 (24)	521	41.5 (61)	
0104	142	86.6 (19)	5.6 (52)	161	44.7 (40)	
0250	174	78.7 (79)	16.7(152)	194	63.9 (1)	
0251	134	67.9(149)	23.9(164)	146	50.7 (15)	
0252	115	80.0 (62)	10.4(116)	156	55.8 (4)	
0253	162	85.8 (24)	4.9 (40)	181	55.8 (4)	
0401	535	89.7 (7)	3.2 (11)	573	38.0 (89)	
0402	498	92.2 (3)	2.6 (4)	539	51.0 (14)	
0403	149	89.9 (6)	4.7 (36)	154	49.4 (21)	
0404	163	85.3 (27)	8.0 (92)	176	39.2 (81)	
0501	552	81.7 (53)	10.7(119)	591	47.2 (26)	
0502	3	* * * *	*	3	*	*
0503	169	75.1(111)	14.2(140)	177	43.5 (45)	
0504	633	85.5 (26)	9.6(107)	687	47.5 (25)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 128  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
0505	461	79.6 (66)	10.4(116)	497	47.7 (24)	
0506	199	75.9(106)	9.0(102)	228	52.2 (9)	
0650	1,076	77.0(100)	11.1(122)	1,144	33.7(117)	
0651	624	79.8 (63)	5.6 (52)	723	45.6 (35)	
0652	397	75.1(111)	14.1(139)	435	39.5 (78)	
0653	443	74.0(120)	14.7(142)	466	44.0 (44)	
0654	372	83.1 (43)	7.0 (80)	405	46.4 (28)	
0655	429	82.1 (50)	9.3(105)	471	39.7 (77)	
0657	118	71.2(138)	12.7(131)	125	46.4 (28)	
0701	430	82.3 (49)	8.8(100)	497	41.0 (66)	
0702	116	88.8 (11)	6.9 (79)	125	49.6 (19)	
0703	254	79.1 (72)	16.1(148)	271	51.3 (13)	
0704	88	* * * *		94	* *	
0705	173	85.0 (31)	5.2 (48)	186	35.5(107)	
0850	1,022	79.3 (69)	9.9(112)	1,199	36.3(104)	
0851	847	79.0 (73)	12.3(126)	928	41.9 (58)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 129  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
0901	1,906	77.2 (97)	10.7(119)	2,119	46.1 (31)	
0902	80	* * * *		83	* *	
1051	1,423	68.1(147)	23.5(163)	1,564	46.4 (28)	
1059	4	* * * *		4	* *	
1060	1	* * * *		1	* *	
1101	1,775	69.4(143)	13.1(135)	2,105	34.2(115)	
1102	888	84.5 (33)	2.6 (4)	990	32.9(119)	
1106	102	75.5(107)	4.9 (40)	117	29.9(129)	
1250	937	70.9(140)	15.5(145)	1,094	40.0 (74)	
1251	827	79.7 (65)	9.7(109)	892	36.1(105)	
1252	819	74.4(118)	13.3(136)	923	37.4 (93)	
1253	773	84.3 (36)	4.8 (38)	856	36.6(102)	
1350	2,575	71.3(136)	16.3(150)	2,799	35.2(108)	
1351	190	77.9 (90)	13.7(137)	205	35.1(109)	
1352	488	63.5(163)	22.1(161)	543	41.1 (64)	
1401	277	88.4 (12)	4.3 (30)	315	51.7 (11)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 130  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
1402	166	72.9(126)	16.3(150)	180	41.1 (64)	
1403	27	* * * *		30	* *	
1404	9	* * * *		21	* *	
1405	0	* * * *		1	* *	
1501	1,085	82.9 (45)	4.7 (36)	1,325	38.9 (83)	
1502	88	* * * *		117	33.3(118)	
1503	1,027	77.5 (94)	6.6 (71)	1,294	36.9 (99)	
1504	2,490	85.8 (24)	2.9 (8)	2,792	36.0(106)	
1505	623	87.3 (15)	4.5 (32)	664	30.4(125)	
1506	453	83.7 (40)	2.4 (3)	550	34.9(111)	
1507	619	91.0 (4)	3.6 (21)	688	40.4 (71)	
1508	143	79.0 (73)	5.6 (52)	280	27.1(137)	
1509	584	78.8 (78)	9.6(107)	713	42.9 (51)	
1510	218	85.3 (27)	3.2 (11)	254	42.9 (51)	
1511	531	85.3 (27)	2.8 (6)	585	38.8 (86)	
1512	309	84.5 (33)	4.5 (32)	322	31.7(123)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 131  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	% (Rank)	% (Rank)
1701	687	76.3(105)	14.4(141)	761	53.9 (6)	
1712	126	71.4(133)	12.7(131)	133	57.9 (3)	
1950	714	74.9(114)	15.5(145)	769	37.2 (96)	
1952	177	81.4 (55)	11.9(125)	181	32.6(121)	
1953	411	81.8 (52)	8.5 (99)	429	36.6(102)	
1954	397	78.6 (80)	12.6(130)	431	43.2 (49)	
1955	212	80.7 (59)	9.4(106)	222	37.8 (90)	
1956	13	* * * *		12	* *	
1957	381	79.5 (68)	8.4 (97)	404	40.6 (68)	
1959	202	77.2 (97)	7.9 (91)	212	26.4(139)	
1963	233	79.0 (73)	12.4(127)	247	41.7 (60)	
2150	729	81.9 (51)	10.6(118)	814	45.7 (33)	
2151	312	60.3(168)	30.4(170)	350	44.3 (41)	
2251	530	80.9 (58)	12.5(128)	595	45.4 (37)	
2252	197	86.8 (17)	4.1 (27)	223	52.9 (7)	
2551	3,998	71.9(130)	7.0 (80)	4,316	25.0(147)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 132  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
2552	991	71.4(133)	8.0 (92)	1,082	21.6(160)	
2554	112	71.4(133)	12.5(128)	122	45.9 (32)	
2555	4,108	78.3 (84)	8.0 (92)	4,354	29.0(132)	
3001	1,044	60.6(167)	18.2(156)	1,226	50.4 (17)	
3002	207	74.4(118)	7.2 (82)	235	51.5 (12)	
3003	22	* * * *		32	* *	
3004	235	53.6(173)	37.9(174)	262	61.8 (2)	
3007	30	* * * *		33	* *	
3009	6	* * * *		8	* *	
3010	0	* * * *		1	* *	
3201	637	89.5 (8)	3.3 (14)	673	43.2 (49)	
3202	19	* * * *		22	* *	
3301	4,123	78.2 (85)	5.5 (51)	4,397	20.9(161)	
3302	439	75.4(108)	3.2 (11)	454	18.5(168)	
3303	79	* * * *		83	* *	
3304	183	83.6 (41)	6.6 (71)	205	40.0 (74)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 133  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
3305	107	86.0 (21)	6.5 (69)	111	25.2(146)	
3306	106	93.4 (1)	2.8 (6)	114	23.7(149)	
3308	67	* * * *		71	* *	
3309	41	* * * *		47	* *	
3311	20	* * * *		19	* *	
3550	96	* * * *		102	47.1 (27)	
3551	33	* * * *		30	* *	
3552	48	* * * *		51	* *	
3650	24	* * * *		33	* *	
3701	736	84.5 (33)	5.7 (55)	792	37.2 (96)	
3702	246	82.5 (47)	6.5 (69)	262	34.4(113)	
3703	117	81.2 (57)	5.1 (43)	123	30.1(127)	
3704	103	77.7 (91)	6.8 (77)	110	28.2(133)	
3805	3,276	83.1 (43)	5.1 (43)	3,616	37.0 (98)	
3950	1,878	77.1 (99)	6.0 (63)	2,256	38.9 (83)	
3952	68	* * * *		79	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 134  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
3953	88	* * * *		100	32.0(122)	
3954	187	84.0 (37)	4.3 (30)	216	30.1(127)	
4001	975	75.0(113)	5.8 (60)	1,164	44.1 (43)	
4002	758	77.4 (95)	4.2 (29)	913	42.9 (51)	
4103	45	* * * *		48	* *	
4106	1,686	61.3(165)	25.6(168)	1,839	37.6 (92)	
4107	3,266	65.4(156)	25.0(167)	3,548	41.2 (63)	
4109	1,081	72.0(129)	19.3(158)	1,119	38.7 (87)	
4110	1,763	54.4(172)	35.4(172)	1,920	49.4 (21)	
4111	62	* * * *		65	* *	
4112	781	43.1(175)	18.1(155)	926	32.8(120)	
4113	820	25.5(177)	51.6(176)	908	40.6 (68)	
4120	2,112	73.2(125)	17.5(154)	2,223	34.7(112)	
4160	2,048	73.3(124)	16.7(152)	2,196	35.0(110)	
4351	706	84.6 (32)	5.7 (55)	779	44.2 (42)	
4352	224	89.3 (9)	3.6 (21)	256	45.7 (33)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 135  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
4353	534	71.3(136)	19.3(158)	603	48.1 (23)	
4354	841	89.3 (9)	3.3 (14)	945	43.5 (45)	
4355	381	65.1(158)	20.7(160)	411	34.3(114)	
4356	585	77.3 (96)	14.0(138)	632	42.6 (54)	
4357	10	*	*	22	*	*
4358	237	50.6(174)	40.9(175)	326	52.1 (10)	
4359	362	79.8 (63)	6.4 (68)	387	41.9 (58)	
4360	211	73.9(121)	16.1(148)	232	40.1 (73)	
4361	258	86.0 (21)	5.4 (50)	284	39.4 (80)	
4362	402	78.6 (80)	11.2(123)	428	43.5 (45)	
4363	893	78.2 (85)	15.0(144)	969	45.0 (39)	
4401	947	56.5(171)	37.6(173)	1,361	38.9 (83)	
4501	1,065	83.9 (39)	6.0 (63)	1,124	40.4 (71)	
4601	1,577	86.7 (18)	3.1 (10)	1,720	26.6(138)	
4701	1,113	79.0 (73)	10.1(114)	1,148	23.7(149)	
5050	932	85.9 (23)	3.3 (14)	1,147	26.2(141)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 136  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
5051	2,440	83.2 (42)	4.0 (24)	2,976	26.0(142)	
5052	1,519	82.5 (47)	3.4 (17)	1,820	26.3(140)	
5054	188	81.4 (55)	7.4 (85)	242	27.7(134)	
5202	342	81.6 (54)	10.8(121)	375	52.5 (8)	
5203	190	85.3 (27)	6.8 (77)	206	49.5 (20)	
5204	154	86.4 (20)	8.4 (97)	172	50.6 (16)	
5205	251	82.9 (45)	10.0(113)	265	42.6 (54)	
5601	1,115	87.0 (16)	4.0 (24)	1,206	40.0 (74)	
5801	2,060	73.9(121)	15.9(147)	2,268	37.8 (90)	
5803	17	* * * *	*	17	* *	
6103	4,469	76.8(101)	11.4(124)	4,847	39.1 (82)	
6301	4,728	75.2(110)	5.7 (55)	5,096	22.8(155)	
6312	2,716	76.5(103)	5.1 (43)	2,909	19.4(164)	
6316	1,107	80.3 (61)	6.6 (71)	1,142	22.9(153)	
6320	261	78.2 (85)	7.3 (83)	267	23.2(151)	
6325	156	78.2 (85)	7.7 (88)	173	26.0(142)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 137  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
6330	280	74.6(116)	5.7 (55)	319	22.6(156)	
6335	612	74.5(117)	8.0 (92)	665	25.7(144)	
6340	538	78.4 (83)	6.7 (75)	586	27.6(136)	
7304	285	79.6 (66)	9.8(110)	311	41.5 (61)	
7306	2,955	70.3(141)	18.4(157)	3,206	42.4 (56)	
7308	215	77.7 (91)	14.9(143)	221	36.7(101)	
7401	2,304	67.0(153)	23.9(164)	2,422	38.3 (88)	
7402	325	84.0 (37)	5.8 (60)	341	25.5(145)	
7403	92	* * * *	*	93	* *	
7404	137	68.6(145)	24.1(166)	140	29.3(131)	
7601	752	78.6 (80)	12.9(134)	810	40.7 (67)	
7850	1,116	34.8(176)	61.1(177)	1,216	50.1 (18)	
7851	214	57.5(170)	33.2(171)	232	43.5 (45)	
7852	125	59.2(169)	29.6(169)	126	37.3 (95)	
8150	89	* * * *	*	109	33.9(116)	
8151	2,622	80.4 (60)	4.8 (38)	2,871	27.7(134)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 138  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
8152	492	78.9 (77)	5.1 (43)	563	45.5 (36)	
8153	31	* * * *		34	* *	
8154	361	92.5 (2)	3.0 (9)	423	37.4 (93)	
8155	72	* * * *		81	* *	
8156	43	* * * *		46	* *	
8157	7	* * * *		9	* *	
8201	2,179	78.2 (85)	6.7 (75)	2,387	19.6(163)	
8202	2,833	73.6(123)	12.7(131)	3,224	36.8(100)	
8253	517	79.3 (69)	4.6 (35)	569	21.8(158)	
8254	607	71.2(138)	9.1(104)	667	18.1(169)	
8255	2,831	77.7 (91)	7.6 (86)	3,100	23.0(152)	
8257	479	71.8(131)	7.3 (83)	517	19.3(165)	
8258	1,612	76.4(104)	8.3 (96)	1,705	22.9(153)	
8901	1,060	65.1(158)	6.1 (65)	1,219	20.3(162)	
8902	557	64.8(160)	7.7 (88)	624	22.6(156)	
8903	1,122	64.3(162)	7.7 (88)	1,297	18.1(169)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 139  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
8904	762	76.6(102)	4.5 (32)	848	23.8(148)	
8905	1,855	75.4(108)	6.3 (67)	2,241	30.4(125)	
9001	74	* * * *		77	* *	
9002	26	* * * *		28	* *	
9004	541	65.4(156)	7.6 (86)	563	11.2(181)	
9005	11	* * * *		11	* *	
9006	30	* * * *		31	* *	
9008	5	* * * *		6	* *	
9009	30	* * * *		34	* *	
9010	264	64.8(160)	6.1 (65)	274	13.1(178)	
9011	14	* * * *		14	* *	
9012	30	* * * *		31	* *	
9014	26	* * * *		26	* *	
9018	10	* * * *		10	* *	
9020	20	* * * *		20	* *	
9022	11	* * * *		12	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 140  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
9023	61	* * * *		65	* *	
9024	29	* * * *		31	* *	
9035	51	* * * *		50	* *	
9039	42	* * * *		45	* *	
9040	431	72.2(128)	2.3 (2)	460	17.4(172)	
9045	1,179	66.8(154)	9.8(110)	1,462	21.7(159)	
9050	1,917	67.4(150)	10.2(115)	2,200	29.5(130)	
9051	2,268	60.8(166)	23.4(162)	2,894	39.5 (78)	
9053	2,771	68.0(148)	5.3 (49)	3,052	18.6(167)	
9054	751	68.3(146)	4.1 (27)	798	16.2(175)	
9058	508	79.3 (69)	8.9(101)	548	40.5 (70)	
9066	2,620	69.3(144)	3.5 (19)	2,927	16.6(174)	
9068	1,555	63.5(163)	5.7 (55)	1,688	17.2(173)	
9069	2,415	71.5(132)	5.0 (42)	2,635	14.1(177)	
9070	2,339	67.1(152)	9.0(102)	2,530	19.1(166)	
9072	1,644	67.2(151)	3.5 (19)	1,803	12.0(180)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6A**  
**2004 - 2006 Pregnancy Nutrition Surveillance <sup>(1)</sup>**  
**Michigan**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Page: 141  
Run Date: 10 / 11 / 07

Clinic	Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>					
	Medical Care			WIC Enrollment		
	1st Trimester	No Care	1st Trimester	Number	%	(Rank)
9074	1,498	65.6(155)	6.6 (71)	1,569	14.6(176)	
9079	1,322	74.7(115)	3.6 (21)	1,363	12.2(179)	
9080	2,105	72.6(127)	5.1 (43)	2,263	30.8(124)	
9151	0	* * * *		1	* *	
Unknown Clinics	322	69.9(142)	5.9 (62)	458	18.1(169)	
<b>Michigan</b>	<b>167,159</b>	<b>74.4</b>	<b>10.6</b>	<b>185,360</b>	<b>32.5</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 142  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household					
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	136	64.0 (176)	121	35.5 (157)	54	* * *	0	* * *	0	* * *	0	* * *
0102	449	58.4 (166)	379	35.4 (156)	176	19.3 (110)	0	* * *	0	* * *	0	* * *
0103	462	60.0 (169)	396	36.6 (161)	180	23.9 (101)	0	* * *	0	* * *	0	* * *
0104	160	62.5 (175)	138	39.1 (168)	71	* * *	0	* * *	0	* * *	0	* * *
0250	195	55.4 (150)	169	30.8 (121)	87	* * *	0	* * *	0	* * *	0	* * *
0251	150	46.7 (92)	130	22.3 (65)	56	* * *	0	* * *	0	* * *	0	* * *
0252	125	52.0 (131)	129	31.0 (122)	46	* * *	0	* * *	0	* * *	0	* * *
0253	171	55.6 (153)	157	34.4 (152)	79	* * *	0	* * *	0	* * *	0	* * *
0401	562	60.1 (170)	497	36.8 (163)	249	30.1 (66)	0	* * *	0	* * *	0	* * *
0402	546	53.1 (141)	470	32.6 (137)	216	29.6 (70)	0	* * *	0	* * *	0	* * *
0403	150	57.3 (162)	138	33.3 (147)	68	* * *	0	* * *	0	* * *	0	* * *
0404	170	55.9 (156)	151	28.5 (107)	62	* * *	0	* * *	0	* * *	0	* * *
0501	585	53.2 (143)	534	29.0 (110)	252	34.9 (35)	0	* * *	0	* * *	0	* * *
0502	3	* * *	3	* * *	1	* * *	0	* * *	0	* * *	0	* * *
0503	175	48.6 (107)	163	24.5 (76)	73	* * *	0	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 143  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0504	659	49.9 (112)	521	27.1 (95)	225	33.3 (39)	0	* * * *	0	* * * *
0505	472	51.9 (129)	411	25.5 (83)	190	37.4 (18)	0	* * * *	0	* * * *
0506	215	51.2 (120)	181	26.0 (90)	81	* * * *	0	* * * *	0	* * * *
0650	1,004	48.3 (103)	886	27.5 (99)	317	31.5 (55)	0	* * * *	0	* * * *
0651	594	72.6 (179)	520	44.6 (173)	258	13.2 (114)	0	* * * *	0	* * * *
0652	372	58.3 (165)	335	32.2 (133)	146	26.0 (92)	0	* * * *	0	* * * *
0653	453	51.7 (125)	427	31.9 (130)	180	26.7 (84)	0	* * * *	0	* * * *
0654	270	80.7 (180)	205	53.7 (174)	105	13.3 (113)	0	* * * *	0	* * * *
0655	459	60.1 (170)	418	38.5 (166)	206	27.7 (75)	0	* * * *	0	* * * *
0657	124	62.1 (174)	107	41.1 (170)	65	* * * *	0	* * * *	0	* * * *
0701	477	47.6 (96)	389	25.2 (81)	148	32.4 (43)	0	* * * *	0	* * * *
0702	119	61.3 (173)	111	32.4 (135)	58	* * * *	0	* * * *	0	* * * *
0703	251	57.0 (161)	244	33.2 (145)	118	38.1 (15)	0	* * * *	0	* * * *
0704	91	* * * *	84	* * * *	41	* * * *	0	* * * *	0	* * * *
0705	178	53.9 (146)	160	34.4 (152)	71	* * * *	0	* * * *	0	* * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 144  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	1,052	52.1 (133)	897	32.9 (142)	306	20.6 (108)	0	* * *	0	* * *
0851	832	50.0 (113)	647	31.7 (129)	218	20.2 (109)	0	* * *	0	* * *
0901	1,976	53.0 (140)	1,764	30.5 (118)	774	28.7 (72)	0	* * *	0	* * *
0902	77	* * *	67	* * *	26	* * *	0	* * *	0	* * *
1051	1,558	47.1 (93)	1,335	25.8 (89)	527	32.3 (45)	0	* * *	0	* * *
1059	4	* * *	3	* * *	1	* * *	0	* * *	0	* * *
1060	1	* * *	1	* * *	0	* * *	0	* * *	0	* * *
1101	2,051	33.3 (44)	1,762	18.8 (51)	466	27.3 (78)	0	* * *	0	* * *
1102	972	47.5 (95)	817	31.6 (128)	348	26.7 (84)	0	* * *	0	* * *
1106	113	50.4 (115)	96	* * *	47	* * *	0	* * *	0	* * *
1250	1,071	42.3 (75)	883	25.5 (83)	288	24.7 (98)	0	* * *	0	* * *
1251	687	67.8 (177)	643	41.5 (171)	337	20.8 (107)	0	* * *	0	* * *
1252	868	34.3 (46)	755	22.8 (68)	197	21.8 (106)	0	* * *	0	* * *
1253	836	46.5 (89)	718	34.8 (154)	277	23.8 (103)	0	* * *	0	* * *
1350	2,715	46.2 (85)	2,385	30.3 (117)	905	25.5 (95)	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 145  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1351	201	40.8 (67)	163	23.9 (74)	61	* * *	0	* * *	0	* * *
1352	539	49.5 (108)	444	32.4 (135)	194	26.3 (90)	0	* * *	0	* * *
1401	313	51.8 (128)	254	33.1 (144)	119	26.9 (82)	0	* * *	0	* * *
1402	181	42.5 (76)	161	16.8 (43)	53	* * *	0	* * *	0	* * *
1403	29	* * *	25	* * *	10	* * *	0	* * *	0	* * *
1404	16	* * *	9	* * *	0	* * *	0	* * *	0	* * *
1405	1	* * *	0	* * *	0	* * *	0	* * *	0	* * *
1501	1,255	37.5 (59)	1,081	19.1 (52)	328	29.9 (68)	0	* * *	0	* * *
1502	107	44.9 (81)	77	* * *	24	* * *	0	* * *	0	* * *
1503	1,118	35.5 (50)	947	23.3 (70)	250	26.8 (83)	0	* * *	0	* * *
1504	2,703	27.7 (33)	2,365	11.6 (26)	513	48.5 (3)	0	* * *	0	* * *
1505	675	37.2 (55)	622	15.4 (40)	189	51.9 (1)	0	* * *	0	* * *
1506	516	49.6 (109)	463	31.5 (127)	172	26.7 (84)	0	* * *	0	* * *
1507	684	47.8 (100)	609	27.3 (97)	254	37.0 (23)	0	* * *	0	* * *
1508	193	0.0 (1)	152	0.0 (1)	0	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 146  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household					
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum			
Clinic	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
1509	652	33.0	(43)	563	20.6	(57)	155	25.2	(97)	0	*	*
1510	248	54.4	(148)	206	33.5	(148)	97	*	*	0	*	*
1511	491	60.3	(172)	455	35.6	(158)	177	26.0	(92)	0	*	*
1512	264	54.2	(147)	241	32.8	(139)	81	*	*	0	*	*
1701	760	51.7	(125)	637	29.7	(113)	295	33.9	(37)	0	*	*
1712	131	57.3	(162)	114	42.1	(172)	61	*	*	0	*	*
1950	755	52.7	(136)	646	32.7	(138)	280	27.1	(80)	0	*	*
1952	171	46.2	(85)	166	28.9	(108)	61	*	*	0	*	*
1953	428	51.6	(123)	390	32.8	(139)	160	30.6	(62)	0	*	*
1954	408	50.7	(118)	361	33.0	(143)	148	28.4	(74)	0	*	*
1955	213	51.6	(123)	192	29.2	(112)	82	*	*	0	*	*
1956	13	*	*	9	*	*	3	*	*	0	*	*
1957	331	56.5	(159)	328	28.4	(105)	124	33.1	(41)	0	*	*
1959	180	58.9	(167)	167	30.5	(118)	56	*	*	0	*	*
1963	231	48.5	(105)	200	25.5	(83)	71	*	*	0	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 147  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)												
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum				
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
2150	776	55.5 (151)	663	28.1 (103)	293	32.1 (51)	0	*	*	0	*	*	
2151	299	59.2 (168)	303	28.1 (103)	135	34.1 (36)	0	*	*	0	*	*	
2251	558	46.6 (90)	497	27.2 (96)	190	31.1 (58)	0	*	*	0	*	*	
2252	218	52.8 (137)	193	32.1 (132)	76	*	*	0	*	*	0	*	*
2551	4,010	36.4 (52)	3,619	22.1 (64)	1,008	31.3 (57)	0	*	*	0	*	*	
2552	1,024	37.2 (55)	856	23.5 (71)	244	31.6 (54)	0	*	*	0	*	*	
2554	122	43.4 (78)	100	26.0 (90)	34	*	*	0	*	*	0	*	*
2555	3,799	50.6 (117)	3,278	30.2 (116)	1,194	26.6 (87)	0	*	*	0	*	*	
3001	1,183	34.7 (47)	1,000	21.7 (63)	290	26.6 (87)	0	*	*	0	*	*	
3002	226	27.9 (35)	201	17.9 (45)	50	*	*	0	*	*	0	*	*
3003	33	*	22	*	1	*	*	0	*	*	0	*	*
3004	261	2.7 (3)	224	0.4 (3)	6	*	*	0	*	*	0	*	*
3007	32	*	28	*	1	*	*	0	*	*	0	*	*
3009	10	*	6	*	0	*	*	0	*	*	0	*	*
3010	1	*	0	*	0	*	*	0	*	*	0	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 148  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3201	666	43.5 (79)	587	23.7 (72)	225	33.3 (39)	0	* * *	0	* * *
3202	22	* *	19	* *	10	* * *	0	* * *	0	* * *
3301	4,026	40.9 (68)	3,485	23.1 (69)	992	32.2 (49)	0	* * *	0	* * *
3302	420	35.5 (50)	387	20.9 (59)	89	* * *	0	* * *	0	* * *
3303	78	* *	68	* *	30	* * *	0	* * *	0	* * *
3304	186	5.4 (5)	168	0.6 (4)	6	* * *	0	* * *	0	* * *
3305	108	35.2 (48)	102	15.7 (41)	23	* * *	0	* * *	0	* * *
3306	111	37.8 (60)	98	* *	22	* * *	0	* * *	0	* * *
3308	69	* *	59	* *	21	* * *	0	* * *	0	* * *
3309	41	* *	37	* *	16	* * *	0	* * *	0	* * *
3311	18	* *	18	* *	7	* * *	0	* * *	0	* * *
3550	103	71.8 (178)	90	* *	56	* * *	0	* * *	0	* * *
3551	32	* *	30	* *	15	* * *	0	* * *	0	* * *
3552	52	* *	46	* *	26	* * *	0	* * *	0	* * *
3650	32	* *	15	* *	5	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 149  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3701	806	45.2 (83)	704	29.0 (110)	273	26.0 (92)	0	* * *	0	* * *
3702	268	53.7 (145)	231	35.1 (155)	99	* * *	0	* * *	0	* * *
3703	125	48.0 (102)	114	28.9 (108)	44	* * *	0	* * *	0	* * *
3704	108	51.9 (129)	98	* * *	40	* * *	0	* * *	0	* * *
3805	3,554	53.4 (144)	3,071	33.7 (150)	1,342	26.5 (89)	0	* * *	0	* * *
3950	1,879	43.3 (77)	1,827	26.8 (94)	582	26.1 (91)	0	* * *	0	* * *
3952	62	* * *	58	* * *	16	* * *	0	* * *	0	* * *
3953	86	* * *	84	* * *	26	* * *	0	* * *	0	* * *
3954	186	39.8 (64)	188	16.5 (42)	56	* * *	0	* * *	0	* * *
4001	1,128	37.0 (53)	875	23.7 (72)	260	24.6 (99)	0	* * *	0	* * *
4002	868	47.7 (99)	700	31.9 (130)	276	24.3 (100)	0	* * *	0	* * *
4103	44	* * *	42	* * *	19	* * *	0	* * *	0	* * *
4106	1,818	24.6 (29)	1,600	13.4 (31)	311	33.4 (38)	0	* * *	0	* * *
4107	3,417	40.2 (66)	2,887	19.6 (54)	930	37.7 (16)	0	* * *	0	* * *
4109	1,136	46.3 (87)	1,024	27.5 (99)	372	30.6 (62)	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 150  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4110	1,900	29.8 (38)	1,655	17.3 (44)	405	30.4 (64)	0	* * *	0	* * *
4111	62	* *	59	* *	24	* * *	0	* * *	0	* * *
4112	907	1.5 (2)	818	0.1 (2)	12	* * *	0	* * *	0	* * *
4113	842	13.1 (9)	757	5.4 (9)	73	* * *	0	* * *	0	* * *
4120	2,204	39.2 (63)	1,997	20.2 (56)	603	37.1 (22)	0	* * *	0	* * *
4160	2,097	30.3 (39)	1,850	15.1 (39)	446	41.0 (12)	0	* * *	0	* * *
4351	762	52.8 (137)	671	33.5 (148)	310	27.7 (75)	0	* * *	0	* * *
4352	241	56.4 (157)	193	37.3 (164)	102	25.5 (95)	0	* * *	0	* * *
4353	592	45.1 (82)	517	25.0 (80)	209	36.8 (26)	0	* * *	0	* * *
4354	930	49.6 (109)	819	30.6 (120)	349	31.8 (53)	0	* * *	0	* * *
4355	401	57.6 (164)	365	33.2 (145)	166	30.7 (61)	0	* * *	0	* * *
4356	620	47.6 (96)	552	24.8 (79)	221	37.6 (17)	0	* * *	0	* * *
4357	15	* *	10	* *	0	* * *	0	* * *	0	* * *
4358	281	3.9 (4)	245	1.2 (5)	8	* * *	0	* * *	0	* * *
4359	382	37.2 (55)	335	25.7 (88)	104	29.8 (69)	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 151  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household					
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4360	235	48.5 (105)	205	32.2 (133)	81	* * * * * *	0	* * * * * *	0	* * * * * *	0	* * * * * *
4361	274	55.8 (155)	249	38.6 (167)	120	22.5 (105)	0	* * * * * *	0	* * * * * *	0	* * * * * *
4362	429	56.4 (157)	398	38.4 (165)	192	27.1 (80)	0	* * * * * *	0	* * * * * *	0	* * * * * *
4363	957	52.1 (133)	856	31.3 (125)	389	30.8 (60)	0	* * * * * *	0	* * * * * *	0	* * * * * *
4401	725	49.8 (111)	1,185	30.0 (114)	229	30.1 (66)	0	* * * * * *	0	* * * * * *	0	* * * * * *
4501	1,124	50.5 (116)	1,030	27.4 (98)	431	32.5 (42)	0	* * * * * *	0	* * * * * *	0	* * * * * *
4601	1,694	44.5 (80)	1,511	25.6 (86)	549	36.4 (27)	0	* * * * * *	0	* * * * * *	0	* * * * * *
4701	1,110	51.4 (122)	1,024	27.7 (101)	404	37.4 (18)	0	* * * * * *	0	* * * * * *	0	* * * * * *
5050	1,056	46.4 (88)	263	14.1 (34)	65	* * * * * *	0	* * * * * *	0	* * * * * *	0	* * * * * *
5051	2,696	41.2 (69)	831	15.0 (38)	176	41.5 (11)	0	* * * * * *	0	* * * * * *	0	* * * * * *
5052	1,670	39.0 (62)	441	14.1 (34)	85	* * * * * *	0	* * * * * *	0	* * * * * *	0	* * * * * *
5054	221	24.0 (28)	100	7.0 (13)	10	* * * * * *	0	* * * * * *	0	* * * * * *	0	* * * * * *
5202	365	46.6 (90)	331	26.6 (92)	132	31.1 (58)	0	* * * * * *	0	* * * * * *	0	* * * * * *
5203	200	55.5 (151)	183	36.1 (159)	83	* * * * * *	0	* * * * * *	0	* * * * * *	0	* * * * * *
5204	170	47.6 (96)	146	21.2 (60)	61	* * * * * *	0	* * * * * *	0	* * * * * *	0	* * * * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 152  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5205	255	54.5 (149)	235	32.8 (139)	93	* * *	0	* * *	0	* * *
5601	1,178	48.4 (104)	1,039	24.7 (78)	409	36.4 (27)	0	* * *	0	* * *
5801	2,261	52.1 (133)	1,935	31.3 (125)	810	32.2 (49)	0	* * *	0	* * *
5803	17	* * *	14	* * *	5	* * *	0	* * *	0	* * *
6103	4,334	47.3 (94)	4,074	26.7 (93)	1,416	27.5 (77)	0	* * *	0	* * *
6301	4,822	31.2 (40)	4,385	18.2 (46)	1,011	35.0 (34)	0	* * *	0	* * *
6312	2,850	23.1 (24)	2,553	11.8 (27)	439	43.1 (7)	0	* * *	0	* * *
6316	1,153	42.0 (73)	1,071	24.6 (77)	328	35.1 (33)	0	* * *	0	* * *
6320	255	22.0 (20)	189	13.8 (33)	23	* * *	0	* * *	0	* * *
6325	164	51.2 (120)	125	39.2 (169)	60	* * *	0	* * *	0	* * *
6330	292	37.3 (58)	205	21.5 (62)	50	* * *	0	* * *	0	* * *
6335	638	37.0 (53)	495	19.2 (53)	127	37.0 (23)	0	* * *	0	* * *
6340	520	39.8 (64)	393	22.6 (67)	106	30.2 (65)	0	* * *	0	* * *
7304	312	38.1 (61)	242	25.6 (86)	75	* * *	0	* * *	0	* * *
7306	3,174	27.7 (33)	2,427	19.7 (55)	566	35.3 (31)	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 153  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7308	224	35.3 (49)	187	21.4 (61)	52	* * * * * *	0	* * * * * *	0	* * * * * *
7401	2,442	55.6 (153)	2,152	34.3 (151)	1,061	28.9 (71)	0	* * * * * *	0	* * * * * *
7402	350	51.7 (125)	311	31.2 (124)	130	32.3 (45)	0	* * * * * *	0	* * * * * *
7403	93	* * * *	86	* * * *	37	* * * * * *	0	* * * * * *	0	* * * * * *
7404	144	56.9 (160)	132	28.0 (102)	69	* * * * * *	0	* * * * * *	0	* * * * * *
7601	808	51.0 (119)	746	28.4 (105)	323	35.3 (31)	0	* * * * * *	0	* * * * * *
7850	535	52.0 (131)	1,080	36.6 (161)	170	18.2 (111)	0	* * * * * *	0	* * * * * *
7851	194	53.1 (141)	205	36.1 (159)	75	* * * * * *	0	* * * * * *	0	* * * * * *
7852	108	52.8 (137)	113	31.0 (122)	43	* * * * * *	0	* * * * * *	0	* * * * * *
8150	101	41.6 (71)	81	* * * *	24	* * * * * *	0	* * * * * *	0	* * * * * *
8151	2,615	32.4 (42)	2,303	18.7 (50)	500	32.4 (43)	0	* * * * * *	0	* * * * * *
8152	546	45.8 (84)	478	22.4 (66)	184	39.1 (13)	0	* * * * * *	0	* * * * * *
8153	31	* * * *	29	* * * *	11	* * * * * *	0	* * * * * *	0	* * * * * *
8154	385	9.9 (7)	321	3.7 (7)	24	* * * * * *	0	* * * * * *	0	* * * * * *
8155	69	* * * *	74	* * * *	4	* * * * * *	0	* * * * * *	0	* * * * * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 154  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8156	44	* * *	41	* * *	18	* * *	0	* * *	0	* * *
8157	7	* * *	5	* * *	0	* * *	0	* * *	0	* * *
8201	2,216	41.2 (69)	1,753	20.7 (58)	529	32.3 (45)	0	* * *	0	* * *
8202	2,383	12.9 (8)	2,243	4.8 (8)	174	37.4 (18)	0	* * *	0	* * *
8253	511	41.9 (72)	430	18.4 (47)	103	23.3 (104)	0	* * *	0	* * *
8254	566	47.9 (101)	413	30.0 (114)	138	23.9 (101)	0	* * *	0	* * *
8255	2,309	50.2 (114)	2,037	24.1 (75)	616	32.3 (45)	0	* * *	0	* * *
8257	218	32.1 (41)	199	13.6 (32)	23	* * *	0	* * *	0	* * *
8258	1,558	42.2 (74)	1,325	25.4 (82)	372	27.2 (79)	0	* * *	0	* * *
8901	1,097	22.2 (22)	982	11.4 (25)	136	39.0 (14)	0	* * *	0	* * *
8902	596	22.0 (20)	509	11.2 (23)	83	* * *	0	* * *	0	* * *
8903	1,148	17.9 (17)	1,056	10.6 (21)	122	32.0 (52)	0	* * *	0	* * *
8904	811	14.1 (12)	716	7.1 (14)	78	* * *	0	* * *	0	* * *
8905	1,970	14.3 (13)	1,818	7.8 (15)	185	28.6 (73)	0	* * *	0	* * *
9001	62	* * *	68	* * *	8	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 155  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9002	24	* * *	20	* * *	3	* * *	0	* * *	0	* * *
9004	520	21.0 (19)	493	11.2 (23)	56	* * *	0	* * *	0	* * *
9005	11	* * *	11	* * *	4	* * *	0	* * *	0	* * *
9006	28	* * *	24	* * *	2	* * *	0	* * *	0	* * *
9008	5	* * *	4	* * *	0	* * *	0	* * *	0	* * *
9009	27	* * *	28	* * *	3	* * *	0	* * *	0	* * *
9010	259	22.8 (23)	242	12.8 (29)	37	* * *	0	* * *	0	* * *
9011	12	* * *	11	* * *	0	* * *	0	* * *	0	* * *
9012	31	* * *	28	* * *	5	* * *	0	* * *	0	* * *
9014	23	* * *	23	* * *	3	* * *	0	* * *	0	* * *
9018	10	* * *	9	* * *	3	* * *	0	* * *	0	* * *
9020	18	* * *	18	* * *	1	* * *	0	* * *	0	* * *
9022	8	* * *	8	* * *	1	* * *	0	* * *	0	* * *
9023	55	* * *	54	* * *	5	* * *	0	* * *	0	* * *
9024	28	* * *	27	* * *	2	* * *	0	* * *	0	* * *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 156  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household					
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum			
Clinic	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
9035	42	*	*	41	*	*	7	*	*	0	*	*
9039	36	*	*	35	*	*	4	*	*	0	*	*
9040	442	25.8	(32)	402	12.9	(30)	86	*	*	0	*	*
9045	1,206	13.9	(11)	1,114	6.7	(12)	103	41.7	(10)	0	*	*
9050	2,031	15.1	(14)	1,912	8.2	(16)	200	37.0	(23)	0	*	*
9051	1,446	16.3	(15)	1,452	6.2	(11)	107	44.9	(5)	0	*	*
9053	2,446	25.3	(31)	1,984	14.9	(36)	308	31.5	(55)	0	*	*
9054	793	23.6	(26)	698	14.9	(36)	101	35.6	(29)	0	*	*
9058	454	6.4	(6)	457	3.1	(6)	19	*	*	0	*	*
9066	1,259	34.2	(45)	1,387	18.5	(48)	144	15.3	(112)	0	*	*
9068	1,155	29.7	(37)	1,098	12.6	(28)	161	37.3	(21)	0	*	*
9069	2,285	23.6	(26)	1,934	10.5	(20)	297	44.4	(6)	0	*	*
9070	1,807	23.5	(25)	1,755	10.0	(18)	202	35.6	(29)	0	*	*
9072	1,759	20.8	(18)	1,567	8.6	(17)	217	41.9	(9)	0	*	*
9074	1,503	25.1	(30)	1,311	11.1	(22)	205	45.4	(4)	0	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Page: 157  
Run Date: 10 / 11 / 07

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	%	Number	%	Number	%	Number	%	Number	%		
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)		
9079	1,331	13.2 (10)	1,183	5.4 (9)	109	50.5 (2)	0	*	*	0	*	*
9080	2,244	17.4 (16)	1,855	10.0 (18)	211	42.7 (8)	0	*	*	0	*	*
9151	1	*	0	*	0	*	0	*	*	0	*	*
Unknown Clinics	346	29.5 (36)	260	18.5 (48)	51	*	0	*	*	0	*	*
<b>Michigan</b>	<b>167,571</b>	<b>38.4</b>	<b>145,412</b>	<b>21.9</b>	<b>41,831</b>	<b>31.3</b>	<b>0</b>	<b>*</b>		<b>0</b>	<b>*</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 158  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number % (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)			
0101	92 * * * *		133 0.0 (1)	92 * *		131	55.7 (88)			
0102	330 6.7 (78)	16.4 (170)	413 0.0 (1)	329 6.7 (78)	399	42.4 (147)				
0103	410 7.3 (107)	6.6 (40)	468 0.2 (121)	409 7.1 (100)	460	43.9 (143)				
0104	127 6.3 (69)	9.4 (107)	149 0.0 (1)	126 6.3 (68)	143	45.5 (137)				
0250	164 4.9 (26)	9.1 (98)	167 0.0 (1)	164 4.9 (26)	159	49.1 (126)				
0251	126 4.0 (10)	5.6 (27)	129 0.8 (170)	125 4.0 (10)	127	48.0 (132)				
0252	132 5.3 (39)	8.3 (76)	133 0.0 (1)	131 5.3 (39)	124	64.5 (30)				
0253	154 7.8 (123)	12.3 (159)	156 0.0 (1)	154 7.8 (122)	147	56.5 (78)				
0401	498 9.0 (143)	10.4 (131)	502 0.0 (1)	498 9.0 (144)	482	52.3 (108)				
0402	475 4.0 (10)	10.1 (124)	474 0.0 (1)	473 4.0 (10)	466	67.8 (24)				
0403	144 6.9 (88)	13.2 (166)	144 0.0 (1)	143 7.0 (94)	139	51.8 (110)				
0404	149 2.0 (1)	11.4 (152)	156 0.0 (1)	149 2.0 (1)	149	51.7 (111)				
0501	518 6.2 (63)	10.6 (134)	548 0.0 (1)	514 6.2 (62)	481	71.1 (19)				
0502	3 * * * *		3 * *	3 * *	3	* *				

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 159  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)								
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0503	156	7.7(120)	9.0 (93)	171	0.0 (1)	156	7.7(118)	167	77.2 (13)
0504	571	7.0 (95)	9.8 (119)	614	0.0 (1)	571	7.0 (94)	555	79.3 (10)
0505	420	6.0 (58)	11.2 (149)	450	0.0 (1)	420	6.0 (58)	405	74.6 (14)
0506	177	3.4 (7)	9.0 (93)	189	0.0 (1)	177	3.4 (7)	182	69.2 (22)
0650	1,001	6.5 (75)	9.4 (107)	1,046	0.0 (1)	998	6.5 (76)	986	69.3 (21)
0651	589	6.5 (75)	6.5 (39)	630	0.3 (145)	587	6.5 (76)	563	50.8(118)
0652	373	5.4 (44)	9.4 (107)	383	0.0 (1)	371	5.4 (44)	366	54.4 (96)
0653	397	4.8 (21)	8.8 (89)	430	0.2 (121)	394	4.8 (21)	421	58.7 (63)
0654	305	8.5(136)	16.7 (171)	320	0.0 (1)	304	8.6(137)	298	63.1 (37)
0655	421	4.8 (21)	8.8 (89)	424	0.0 (1)	421	4.8 (21)	406	51.5(113)
0657	105	6.7 (78)	9.5 (112)	112	0.0 (1)	105	6.7 (78)	111	63.1 (37)
0701	359	2.8 (3)	10.3 (127)	411	0.2 (121)	359	2.8 (3)	397	74.6 (14)
0702	104	4.8(21)	12.5 (162)	110	0.0 (1)	104	4.8 (21)	109	46.8(135)
0703	233	4.3 (15)	10.7 (139)	235	0.0 (1)	233	4.3 (15)	212	56.1 (82)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 160  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)	Number (Rank)			
0704	81	* * * *	82	*	*	80	*	*	79	* *
0705	149	4.0 (10) 9.4 (107)	170	0.0 (1)		149	4.0 (10)		164	58.5 (64)
0850	969	7.6(118) 9.9 (121)	1,044	0.1 (99)		963	7.7(118)		1,000	61.5 (44)
0851	808	6.8 (84) 11.0 (144)	839	0.1 (99)		805	6.8 (84)		819	52.5(107)
0901	1,827	5.5 (48) 10.6 (134)	1,874	0.1 (99)		1,823	5.5 (47)		1,822	50.5(120)
0902	70	* * * *	71	*	*	70	*	*	70	* *
1051	1,342	7.2(103) 10.7 (139)	1,386	0.1 (99)		1,338	7.2(105)		1,338	78.6 (11)
1059	3	* * * *	4	*	*	3	*	*	4	* *
1060	1	* * * *	1	*	*	1	*	*	1	* *
1101	1,763	8.1(130) 7.4 (55)	1,773	0.0 (1)		1,756	8.1(130)		1,705	45.6(136)
1102	825	5.9 (54) 8.4 (79)	856	0.1 (99)		822	5.8 (53)		819	51.6(112)
1106	98	* * * *	102	0.0 (1)		98	*	*	96	* *
1250	927	5.0 (32) 9.0 (93)	966	0.0 (1)		922	5.0 (31)		946	57.4 (71)
1251	802	4.7 (19) 7.6 (59)	820	0.0 (1)		802	4.7 (19)		771	56.2 (80)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 161  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)	Number (Rank)			
1252	698	5.3 (39) 7.2 (52)	809	0.4 (157)	696	5.3 (39)	777	54.1 (100)		
1253	720	8.8(139) 7.1 (51)	744	0.1 (99)	716	8.8(139)	709	42.6(146)		
1350	2,384	6.9 (88) 9.6 (114)	2,515	0.0 (1)	2,380	6.9 (88)	2,276	48.6(127)		
1351	183	4.9 (26) 12.6 (163)	187	0.0 (1)	182	4.9 (26)	179	59.2 (61)		
1352	450	6.7 (78) 7.8 (66)	463	0.4 (157)	447	6.7 (78)	443	34.1(163)		
1401	254	7.1 (99) 9.8 (119)	268	0.0 (1)	253	7.1(100)	259	71.8 (17)		
1402	169	3.0 (5) 17.2 (172)	173	0.0 (1)	168	3.0 (5)	169	85.8 (9)		
1403	26	* * * *	27	* *	26	* *	25	* *		
1404	10	* * * *	11	* *	10	* *	11	* *		
1405	0	* * * *	0	* *	0	* *	0	* *		
1501	1,026	5.0 (32) 9.2 (102)	1,132	0.0 (1)	1,020	5.0 (31)	1,088	59.4 (59)		
1502	87	* * * *	88	* *	87	* *	78	* *		
1503	1,020	7.4(111) 9.6 (114)	1,115	0.0 (1)	1,018	7.4(111)	1,025	62.4 (41)		
1504	2,117	7.8(123) 8.5 (83)	2,478	0.1 (99)	2,112	7.8(122)	2,377	65.2 (29)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 162  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)							
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1505	594	6.7 (78) 10.9 (142)	625	0.2 (121)	592	6.8 (84)	605	64.0 (33)
1506	453	4.2 (13) 12.1 (157)	477	0.2 (121)	452	4.2 (13)	467	55.9 (84)
1507	548	4.9 (26) 12.0 (156)	607	0.0 (1)	545	5.0 (31)	600	59.8 (57)
1508	146	6.2 (63) 12.3 (159)	199	0.0 (1)	145	6.2 (62)	193	89.6 (7)
1509	428	5.8 (53) 8.2 (73)	599	0.2 (121)	427	5.9 (55)	580	61.7 (43)
1510	155	5.2 (38) 9.0 (93)	221	0.5 (163)	153	5.2 (38)	216	55.6 (89)
1511	508	5.3 (39) 11.2 (149)	525	0.0 (1)	508	5.3 (39)	516	56.2 (80)
1512	280	7.9 (128) 7.9 (70)	292	0.0 (1)	280	7.9 (128)	286	58.0 (66)
1701	654	4.4 (16) 13.9 (168)	661	0.0 (1)	654	4.4 (16)	642	63.2 (36)
1712	119	3.4 (7) 11.8 (154)	124	0.0 (1)	118	3.4 (7)	118	55.9 (84)
1950	672	6.1 (59) 7.0 (49)	681	0.1 (99)	667	6.1 (59)	634	56.6 (76)
1952	171	11.1 (154) 9.9 (121)	172	0.6 (166)	169	11.2 (155)	162	58.0 (66)
1953	398	7.5 (114) 9.5 (112)	403	0.7 (169)	391	7.7 (118)	383	52.7 (106)
1954	343	7.6 (118) 8.7 (87)	374	0.3 (145)	337	7.4 (111)	347	55.6 (89)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 163  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)	Number (Rank)			
1955	186	5.4 (44) 8.6 (86)	196	0.5 (163)	183	5.5 (47)	188	59.0 (62)		
1956	11	* * * *	11	* *	11	* *	11	*	*	
1957	347	6.1 (59) 11.0 (144)	370	0.0 (1)	347	6.1 (59)	350	59.7 (58)		
1959	178	6.2 (63) 6.7 (44)	197	0.0 (1)	178	6.2 (62)	182	59.3 (60)		
1963	194	5.7 (52) 6.7 (44)	212	0.0 (1)	192	5.7 (52)	200	60.0 (54)		
2150	687	6.8 (84) 11.8 (154)	703	0.1 (99)	686	6.7 (78)	644	60.1 (53)		
2151	310	4.8 (21) 7.4 (55)	317	0.3 (145)	310	4.8 (21)	280	45.0 (138)		
2251	520	5.4 (44) 10.6 (134)	530	0.4 (157)	518	5.4 (44)	525	57.9 (68)		
2252	190	2.1 (2) 11.1 (147)	193	0.0 (1)	189	2.1 (2)	188	55.9 (84)		
2551	3,718	10.2(151) 4.7 (10)	3,815	0.0 (1)	3,710	10.2(151)	3,600	36.9(158)		
2552	819	8.9(142) 5.4 (19)	922	0.2 (121)	817	8.9(143)	912	43.5(145)		
2554	91	* * * *	103	0.0 (1)	91	* *	97	*	*	
2555	3,582	6.8 (84) 8.4 (79)	3,903	0.2 (121)	3,573	6.9 (88)	3,802	48.4 (129)		
3001	1,004	11.6(158) 6.1 (30)	1,011	0.0 (1)	1,000	11.6(158)	947	32.6(164)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 164  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)										
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed			
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)			
3002	201	10.9(153)	5.0 (12)	202	0.0 (1)	200	11.0(153)	187	28.3(169)		
3003	21	*	*	*	*	21	*	*	21	*	*
3004	225	4.9 (26)	5.3 (18)	226	0.0 (1)	225	4.9 (26)	220	78.6 (11)		
3007	26	*	*	*	*	28	*	*	28	*	*
3009	5	*	*	*	*	5	*	*	5	*	*
3010	0	*	*	*	*	0	*	*	0	*	*
3201	547	5.3 (39)	11.2 (149)	594	0.2 (121)	544	5.3 (39)	575	44.2(141)		
3202	18	*	*	*	*	19	*	*	19	*	*
3301	3,513	7.5(114)	8.4 (79)	3,979	0.2 (121)	3,494	7.5(115)	3,613	61.4 (46)		
3302	388	7.5(114)	7.0 (49)	425	0.0 (1)	388	7.5(115)	377	57.0 (72)		
3303	75	*	*	*	*	79	*	*	77	*	*
3304	163	5.5 (48)	5.5 (23)	180	0.0 (1)	161	5.6 (49)	177	93.2 (5)		
3305	97	*	*	*	*	105	0.0 (1)	97	*	*	
3306	102	8.8(139)	12.7 (165)	111	0.9 (174)	102	8.8(139)	101	64.4 (31)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 165  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (%) (Rank)							
3308	55 * * * *	65 * *	55 * *	60 * *						
3309	34 * * * *	40 * *	33 * *	35 * *						
3311	19 * * * *	20 * *	18 * *	19 * *						
3550	89 * * * *	88 * *	88 * *	88 * *						
3551	30 * * * *	30 * *	30 * *	30 * *						
3552	46 * * * *	45 * *	45 * *	45 * *						
3650	15 * * * *	14 * *	14 * *	14 * *						
3701	702 5.3 (39) 11.1 (147)	702 0.0 (1)	700 5.3 (39)	660 50.8(118)						
3702	233 11.2(156) 7.7 (61)	234 0.0 (1)	233 11.2(155)	218 49.5(123)						
3703	114 7.0 (95) 3.5 (3)	114 0.0 (1)	114 7.0 (94)	110 50.9(116)						
3704	98 * * * *	98 * *	98 * *	88 * *						
3805	3,161 7.7(120) 7.6 (59)	3,227 0.0 (1)	3,154 7.7(118)	3,096 50.3(121)						
3950	1,659 8.4(135) 10.4 (131)	1,973 0.2 (121)	1,655 8.3(133)	1,824 56.7 (75)						
3952	49 * * * *	62 * *	48 * *	58 * *						

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 166  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (%) (Rank)							
3953	82	* * * *	93	* *	82	* *	85	* *		
3954	165	4.2 (13) 12.1 (157)	196	0.0 (1)	165	4.2 (13)	182	61.0 (47)		
4001	841	10.0 (149) 5.5 (23)	907	0.3 (145)	834	10.0 (149)	827	53.6 (103)		
4002	682	8.2 (132) 9.2 (102)	734	0.0 (1)	678	8.3 (133)	698	56.9 (73)		
4103	39	* * * *	40	* *	39	* *	37	* *		
4106	1,594	8.5 (136) 5.5 (23)	1,612	0.0 (1)	1,592	8.5 (136)	1,389	54.6 (94)		
4107	2,929	9.3 (146) 9.1 (98)	3,081	0.1 (99)	2,918	9.2 (146)	2,860	54.4 (96)		
4109	1,025	7.9 (128) 10.6 (134)	1,032	0.0 (1)	1,023	7.9 (128)	967	60.3 (51)		
4110	1,620	5.0 (32) 9.4 (107)	1,651	0.0 (1)	1,618	5.0 (31)	1,571	59.9 (55)		
4111	59	* * * *	59	* *	59	* *	52	* *		
4112	802	4.9 (26) 10.3 (127)	806	0.0 (1)	799	4.8 (21)	772	89.6 (7)		
4113	764	3.0 (5) 9.3 (105)	769	0.0 (1)	762	3.0 (5)	726	71.2 (18)		
4120	2,005	6.2 (63) 9.6 (114)	2,012	0.0 (1)	1,998	6.3 (68)	1,848	57.5 (70)		
4160	1,943	7.1 (99) 9.1 (98)	1,981	0.2 (121)	1,937	7.0 (94)	1,930	55.2 (92)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 167  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number % (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)		
4351	661	5.0 (32) 8.0 (72)	683	0.1 (99)	660	5.0 (31)	635	54.3 (98)		
4352	210	7.1 (99) 11.0 (144)	212	0.0 (1)	210	7.1 (100)	202	52.0 (109)		
4353	523	6.9 (88) 7.8 (66)	529	0.0 (1)	521	6.9 (88)	502	58.4 (65)		
4354	802	6.7 (78) 12.3 (159)	828	0.0 (1)	799	6.8 (84)	810	62.5 (39)		
4355	362	6.9 (88) 7.5 (57)	364	0.0 (1)	362	6.9 (88)	332	62.3 (42)		
4356	550	6.4 (71) 7.8 (66)	556	0.2 (121)	546	6.4 (71)	536	56.0 (83)		
4357	11	* * * *	15	* *	11	* *	15	* *		
4358	242	5.4 (44) 8.3 (76)	256	0.0 (1)	242	5.4 (44)	255	65.5 (28)		
4359	338	6.5 (75) 10.9 (142)	355	0.3 (145)	335	6.3 (68)	322	60.6 (48)		
4360	199	4.5 (18) 10.6 (134)	196	0.0 (1)	196	4.6 (18)	189	64.0 (33)		
4361	250	6.8 (84) 6.0 (29)	251	0.0 (1)	250	6.8 (84)	236	56.8 (74)		
4362	396	7.3 (107) 6.3 (33)	396	0.0 (1)	394	7.4 (111)	378	70.6 (20)		
4363	853	5.9 (54) 11.5 (153)	856	0.0 (1)	847	5.8 (53)	815	63.8 (35)		
4401	1,182	7.2 (103) 8.9 (92)	1,212	0.2 (121)	1,178	7.2 (105)	1,169	49.5 (123)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 168  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)							
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4501	1,017	6.7 (78) 10.0 (123)	1,029	0.1 (99)	1,014	6.7 (78)	1,006	57.6 (69)
4601	1,528	6.3 (69) 6.7 (44)	1,528	0.1 (99)	1,524	6.4 (71)	1,505	54.0 (101)
4701	1,018	5.9 (54) 10.5 (133)	1,058	0.5 (163)	1,009	5.9 (55)	1,008	67.8 (24)
5050	329	8.8 (139) 10.3 (127)	343	0.0 (1)	328	8.8 (139)	343	95.3 (2)
5051	1,031	5.9 (54) 7.7 (61)	1,080	0.3 (145)	1,027	5.9 (55)	1,073	94.5 (3)
5052	514	7.0 (95) 6.4 (36)	532	0.4 (157)	510	7.1 (100)	529	93.6 (4)
5054	107	2.8 (3) 8.4 (79)	124	0.0 (1)	106	2.8 (3)	125	99.2 (1)
5202	316	4.7 (19) 10.8 (141)	329	0.0 (1)	316	4.7 (19)	305	66.6 (27)
5203	179	5.0 (32) 10.1 (124)	184	0.0 (1)	178	5.1 (37)	172	59.9 (55)
5204	144	4.9 (26) 13.9 (168)	148	0.0 (1)	143	4.9 (26)	140	68.6 (23)
5205	234	10.7 (152) 9.0 (93)	238	0.4 (157)	232	10.8 (152)	227	60.4 (49)
5601	1,054	7.2 (103) 9.6 (114)	1,064	0.1 (99)	1,053	7.2 (105)	1,006	62.5 (39)
5801	1,938	7.0 (95) 6.3 (33)	1,967	0.1 (99)	1,925	7.0 (94)	1,876	48.0 (132)
5803	14	* * * *	14	* *	14	* *	13	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 169  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)							
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6103	3,955	7.3(107)	7.7 (61)	4,194	0.0 (1)	3,942	7.3(109)	3,957 41.9(148)
6301	4,469	7.8(123)	7.8 (66)	4,598	0.0 (1)	4,465	7.8(122)	4,446 47.2(134)
6312	2,630	8.3(134)	8.2 (73)	2,680	0.1 (99)	2,621	8.3(133)	2,614 53.9(102)
6316	1,080	7.4(111)	7.9 (70)	1,085	0.0 (1)	1,076	7.4(111)	1,085 56.6 (76)
6320	213	6.1 (59)	10.3 (127)	233	0.9 (174)	210	6.2 (62)	227 55.5 (91)
6325	124	4.8 (21)	9.7 (118)	142	0.0 (1)	123	4.9 (26)	138 49.3(125)
6330	219	12.3(162)	6.8 (48)	244	0.8 (170)	217	12.4(164)	231 37.2(157)
6335	487	6.2 (63)	6.6 (40)	559	0.0 (1)	487	6.2 (62)	551 48.1(131)
6340	422	6.9 (88)	6.6 (40)	482	0.2 (121)	416	6.7 (78)	459 54.5 (95)
7304	270	3.7 (9)	12.6 (163)	272	0.4 (157)	269	3.7 (9)	255 43.9(143)
7306	2,821	8.2(132)	8.7 (87)	2,837	0.1 (99)	2,816	8.1(130)	2,586 41.7(149)
7308	204	4.4 (16)	13.7 (167)	205	0.0 (1)	204	4.4 (16)	189 60.3 (51)
7401	2,146	7.2(103)	6.4 (36)	2,165	0.0 (1)	2,142	7.2(105)	2,124 44.7(140)
7402	306	6.2 (63)	8.8 (89)	310	0.0 (1)	306	6.2 (62)	298 45.0(138)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 170  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)	Number (Rank)			
7403	87	* * * *	88	*	*	87	*	*	86	* *
7404	131	6.9 (88) 1.5 (1)	133	0.8 (170)	131	6.9 (88)	133	51.1 (114)		
7601	748	6.4 (71) 7.5 (57)	752	0.0 (1)	746	6.4 (71)	716	52.8 (105)		
7850	1,051	6.1 (59) 7.3 (54)	1,065	0.0 (1)	1,051	6.1 (59)	989	53.2 (104)		
7851	202	6.4 (71) 6.4 (36)	206	0.0 (1)	202	6.4 (71)	187	60.4 (49)		
7852	109	6.4 (71) 10.1 (124)	113	0.0 (1)	109	6.4 (71)	109	64.2 (32)		
8150	76	* * * *	83	*	*	73	*	*	81	* *
8151	2,316	7.3 (107) 9.1 (98)	2,488	0.2 (121)	2,299	7.0 (94)	2,314	61.5 (44)		
8152	469	8.5 (136) 4.1 (4)	477	0.2 (121)	465	8.6 (137)	444	50.0 (122)		
8153	31	* * * *	32	*	*	31	*	*	32	* *
8154	337	5.0 (32) 7.7 (61)	354	0.0 (1)	337	5.0 (31)	352	91.2 (6)		
8155	69	* * * *	74	*	*	69	*	*	69	* *
8156	38	* * * *	42	*	*	38	*	*	40	* *
8157	6	* * * *	7	*	*	6	*	*	6	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 171  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)								
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
8201	1,790	6.9 (88)	8.2 (73)	1,849	0.2 (121)	1,779	6.9 (88)	1,776 51.0(115)	
8202	2,334	5.5 (48)	7.7 (61)	2,717	0.1 (99)	2,324	5.6 (49)	2,622 73.7 (16)	
8253	474	7.4(111)	9.3 (105)	498	0.6 (166)	468	7.3(109)	454 50.9(116)	
8254	467	7.7(120)	9.2 (102)	508	0.2 (121)	464	7.8(122)	453 38.6(154)	
8255	2,302	7.8(123)	8.5 (83)	2,445	0.3 (145)	2,287	7.8(122)	2,343 44.1(142)	
8257	346	9.2(145)	5.5 (23)	409	1.0 (176)	340	8.8(139)	399 56.4 (79)	
8258	1,458	7.1 (99)	8.5 (83)	1,484	0.1 (99)	1,455	7.1(100)	1,439 41.3(151)	
8901	931	12.5(166)	4.4 (6)	1,052	0.3 (145)	925	12.5(165)	992 29.0(166)	
8902	504	12.3(162)	5.4 (19)	534	0.0 (1)	504	12.3(162)	521 36.1(161)	
8903	1,019	11.8(160)	3.1 (2)	1,100	0.1 (99)	1,014	11.8(160)	1,054 26.4(171)	
8904	663	10.1(150)	6.3 (33)	758	0.1 (99)	660	10.0(149)	732 48.5(128)	
8905	1,732	8.1(130)	7.2 (52)	1,924	0.2 (121)	1,725	8.1(130)	1,851 36.5(160)	
9001	71	*	*	71	*	*	71	*	*
9002	23	*	*	23	*	*	23	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 172  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)										
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed			
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)			
9004	522	12.8(168)	6.1 (30)	526	0.0 (1)	521	12.7(167)	508	21.3(173)		
9005	11	*	*	*	*	11	*	*	11	*	*
9006	28	*	*	*	*	28	*	*	27	*	*
9008	4	*	*	*	*	4	*	*	4	*	*
9009	28	*	*	*	*	30	*	*	29	*	*
9010	229	14.0(172)	4.8 (11)	249	0.0 (1)	228	14.0(172)	244	19.3(174)		
9011	14	*	*	*	*	14	*	*	14	*	*
9012	30	*	*	*	*	30	*	*	29	*	*
9014	25	*	*	*	*	25	*	*	22	*	*
9018	9	*	*	*	*	9	*	*	9	*	*
9020	18	*	*	*	*	18	*	*	18	*	*
9022	10	*	*	*	*	10	*	*	9	*	*
9023	56	*	*	*	*	56	*	*	54	*	*
9024	27	*	*	*	*	29	*	*	29	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** (1)  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 173  
Run Date: 10 / 11 / 07

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)				
9035	45	* * * *	45	*	*	45	*	*	44	* *
9039	40	* * * *	41	*	*	40	*	*	38	* *
9040	414	9.4(147)	4.1 (4)	417	0.0 (1)	413	9.4(147)	410	25.6(172)	
9045	1,254	9.4(147)	5.7 (28)	1,308	0.2 (121)	1,246	9.5(148)	1,263	39.7(152)	
9050	1,825	7.8(123)	6.6 (40)	1,932	0.0 (1)	1,823	7.8(122)	1,870	48.4(129)	
9051	2,144	5.6 (51)	8.3 (76)	2,527	0.2 (121)	2,128	5.6 (49)	2,416	55.9 (84)	
9053	2,420	11.1(154)	5.1 (15)	2,639	0.1 (99)	2,413	11.1(154)	2,130	36.0(162)	
9054	707	13.9(171)	5.0 (12)	711	0.0 (1)	706	13.9(171)	669	29.0(166)	
9058	429	7.5(114)	5.4 (19)	463	0.2 (121)	428	7.5(115)	460	55.2 (92)	
9066	2,377	13.4(170)	4.4 (6)	2,609	0.2 (121)	2,363	13.4(170)	2,458	28.8(168)	
9068	1,450	12.4(164)	6.1 (30)	1,504	0.3 (145)	1,441	12.5(165)	1,086	37.5(156)	
9069	2,254	12.9(169)	5.1 (15)	2,363	0.0 (1)	2,252	12.8(169)	2,165	41.7(149)	
9070	2,014	12.4(164)	4.6 (9)	2,174	0.3 (145)	2,005	12.3(162)	2,020	30.2(165)	
9072	1,589	12.5(166)	5.0 (12)	1,620	0.8 (170)	1,571	12.7(167)	1,601	27.6(170)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8A**  
**2004 - 2006 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Michigan**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Page: 174  
Run Date: 10 / 11 / 07

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9074	1,406	11.9(161)	4.5 (8)	1,420	0.0 (1)	1,406	11.9(161)	1,305	38.1(155)
9079	1,171	11.5(157)	5.4 (19)	1,225	0.3 (145)	1,165	11.5(157)	1,148	39.3(153)
9080	1,928	9.1(144)	6.7 (44)	1,939	0.3 (145)	1,910	9.1(145)	1,924	54.3 (98)
9151	0	*	*	0	*	0	*	0	*
Unknown Clinics	268	11.6(158)	5.2 (17)	308	0.6 (166)	268	11.6(158)	282	36.9(158)
<b>Michigan</b>	<b>150,761</b>	<b>7.8</b>	<b>8.0</b>	<b>159,123</b>	<b>0.1</b>	<b>150,183</b>	<b>7.8</b>	<b>150,569</b>	<b>51.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.